

ARTICLE

Marine resources alter tundra food web dynamics by subsidizing a terrestrial predator on the sea ice

Sean M. Johnson-Bice¹  | Frank B. Baldwin² | Evan S. Richardson³ | James D. Roth¹

¹Department of Biological Sciences,
University of Manitoba, Winnipeg,
Manitoba, Canada

²Canadian Wildlife Service, Environment
and Climate Change Canada, Winnipeg,
Manitoba, Canada

³Wildlife Research Division, Environment
and Climate Change Canada, Winnipeg,
Manitoba, Canada

Correspondence

Sean M. Johnson-Bice
Email: s.johnsonbice@gmail.com

Funding information

Natural Sciences and Engineering
Research Council of Canada; Natural
Resources Canada Polar Continental Shelf
Program; Environment and Climate
Change Canada, Parks Canada; the
University of Manitoba Fieldwork
Support Program; the Churchill Northern
Studies Centre (CNSC) Northern
Research Fund; the Province of Manitoba,
Earth Ranger Foundation; Isdell Family
Foundation; Wildlife Media

Handling Editor: Daniel S. Gruner

Abstract

Predator use of resource subsidies can strengthen top-down effects on prey when predators respond numerically to subsidies. Although allochthonous subsidies are generally transported along natural gradients, consumers can cross ecosystem boundaries to acquire subsidies, thereby linking disparate ecosystems. In coastal Arctic ecosystems, terrestrial predators like Arctic foxes (*Vulpes lagopus*) cross into the marine environment (sea ice) during winter to access marine resources. Arctic foxes kill seal pups and scavenge seal carrion (often remains from polar bear *Ursus maritimus* kills), especially when rodent abundance is low. Terrestrial predator use of marine subsidies may strengthen the top-down control of tundra food webs, but this hypothesis remained untested. We evaluated tundra food web dynamics at the terrestrial–marine interface from an ecosystem-level perspective by assessing: (1) how winter environmental conditions affect rodent abundance and marine subsidy availability, (2) the response of the Arctic fox population to this seasonal food variability, and (3) the subsequent effects of Arctic foxes on Canada goose (*Branta canadensis interior*) reproduction. Arctic foxes responded numerically to rodent abundance, which was positively related to snow persistence. Arctic fox abundance was positively related to polar bear body condition metrics, which were used as a proxy for marine subsidy availability. Canada goose reproductive success, in turn, was negatively related to Arctic fox abundance. Long-term trends in goose reproduction and snow persistence also indicate an ongoing phenological mismatch between nesting initiation and spring onset. Our results reveal near-term apparent competition between rodents and geese through a shared predator, Arctic foxes, contrasting with prior studies evaluating rodent–goose–predator relationships. Moreover, we establish a link between tundra and sea ice food webs by demonstrating how seal availability can affect goose reproduction indirectly by increasing Arctic fox predation on goose nests via a population response of foxes to marine resources. These marine resources are often provisioned by polar bears, and with both Arctic

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2025 The Author(s). *Ecology* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

foxes and polar bears undergoing long-term regional declines evidently driven by climate-related changes in prey abundance and availability, we contextualize our study within ongoing climate change and highlight the vulnerability of this likely widespread terrestrial–marine linkage in a warming Arctic.

KEYWORDS

allochthonous resources, apparent competition, Arctic fox, Canada goose, carrion provisioning, climate change, indirect effects, lemmings, phenological mismatch, polar bear, predator–prey dynamics, resource subsidy

INTRODUCTION

The flow of allochthonous resources from one ecosystem to another is ubiquitous and can have powerful effects on food webs in recipient ecosystems (Polis et al., 1997). In general, the effects of allochthonous subsidies on recipient food webs are higher when they subsidize systems with low levels of comparable resources (Marczak et al., 2007). Resource subsidies can also have context-dependent effects on consumers depending on the timing, duration, or nutritional content of the subsidies (Subalusky & Post, 2019). For instance, predators may respond numerically to resource subsidies only when they are available at particular times of year (Sato et al., 2016). Subsidies that are available to predators at irregular intervals or only for short periods of time can have short-term positive indirect effects on other prey by reducing predation pressure through consumer diet switching (Spiller et al., 2010). In contrast, spatial subsidies that are regularly available tend to elicit a numerical response in predator populations (Eide et al., 2012; Eriksson et al., 2022; Rose & Polis, 1998), which, in turn, may strengthen top-down effects on other prey through increased predator abundance (i.e., apparent competition; Holt, 1977; Polis et al., 1997).

Generally, allochthonous resources are transported with natural gradients (e.g., wind, current, gravity), but mobile consumers can cross ecosystem boundaries against natural gradients to consume resources, effectively linking disparate ecosystems through their movement (Lundberg & Moberg, 2003; McInturf et al., 2019). Resource dynamics in one ecosystem can indirectly affect another ecosystem via these mobile consumers, including modifying trophic interactions with other prey species. Notable examples of this phenomenon come from the North American Pacific Northwest: wolf (*Canis lupus*) consumption of sea otters (*Enhydra lutris*) and spawning salmon in aquatic ecosystems can subsidize wolf populations and lead to top-down limitation of ungulates through the wolves' numerical response to these marine subsidies (Adams et al., 2010; Roffler et al., 2023). Despite

the recognition that (1) allochthonous subsidies can have strong effects on predator populations, and (2) predators regularly cross ecosystem boundaries to acquire resources, examples that demonstrate resource subsidies from a donor ecosystem can indirectly alter food web dynamics in a recipient ecosystem via mobile predators crossing ecosystem boundaries are lacking, in large part due to the challenges of studying these trophic relationships at relevant spatial and temporal scales.

Despite simple food web structures, complex indirect trophic interactions can arise in tundra ecosystems through fluctuating prey populations and resource availability. Tundra rodents (especially lemmings [*Lemmus* spp. and *Dicrostonyx* spp.]) are renowned for their 3- to 5-year cycles in abundance (Stenseth, 1999), and their irruptive population dynamics can have strong effects on food webs (Gilg et al., 2003, 2006; Schmidt et al., 2012). However, recent work found high variation in the presence, strength, and regularity of lemming cycles (Gauthier et al., 2024), suggesting lemming irruptions are likely driven by both intrinsic (density dependence) and extrinsic factors (food supply, environmental conditions, predator abundance) (Oli, 2019). In particular, snow conditions affect lemming abundance in some systems (Bilodeau et al., 2013a; Kausrud et al., 2008), as winter reproduction plays a major role in lemming population changes (Duchesne et al., 2011; Fauteux et al., 2015). Predators such as Arctic foxes (*Vulpes lagopus*) benefit from high lemming abundance, and their population changes are generally strongly linked to lemming dynamics when available (Angerbjörn et al., 1999; Gilg et al., 2006). However, as opportunistic predators, Arctic foxes readily switch to other foods when rodent abundance is low, which may generate indirect trophic interactions between rodents and other prey via their shared predator (foxes) (Bêty et al., 2001, 2002; Juhasz et al., 2020). For tundra-nesting birds, such as many goose species, predation rates by foxes tend to increase during years of low lemming abundance (Dulude-de Broin et al., 2023; McKinnon et al., 2013). Although Arctic fox diets are influenced by goose presence during summer, changes in

fox abundance are driven more by rodents than by geese where both prey are available (Chevallier et al., 2020; McDonald et al., 2017; Samelius & Alisauskas, 2017).

Ecological theory suggests that ecosystems with a high perimeter to area ratio—like most coastal tundra ecosystems—are more likely to be affected by allochthonous subsidies (Polis et al., 1997), and indeed researchers have found tundra predator diets are often influenced by marine resources (Carbonell Ellgutter et al., 2020; Tarroux et al., 2012; Therrien et al., 2011). During winter, mobile tundra predators like Arctic foxes can easily venture onto the sea ice to access marine subsidies (Lai et al., 2017; Roth, 2002; Warret Rodrigues & Roth, 2023). Arctic foxes frequently scavenge seal carrion provisioned by polar bears (*Ursus maritimus*), which often only consume seal blubber and abandon the rest of the carcass (Stirling & Archibald, 1977). Arctic foxes are also capable of killing seal pups within their lairs (Lydersen & Gjertz, 1986), with one study estimating Arctic foxes killed ~26% of ringed seal (*Pusa hispida*) pups in the area (Smith, 1976). Marine resource usage helps to stabilize Arctic fox population dynamics (Nater et al., 2021), particularly when other foods are scarce (Roth, 2003). Yet, with the Arctic warming four times faster than the rest of the world (Rantanen et al., 2022), changing climatic conditions may alter the availability of resource subsidies to tundra predators. On the one hand, rain-on-snow events, which are occurring more frequently in many Arctic ecosystems, can facilitate predator access to terrestrial carrion by increasing reindeer (*Rangifer tarandus*) mortality (Ehrich et al., 2017; Hansen et al., 2013). On the other hand, degrading sea ice regimes may reduce tundra predator access to marine subsidies in winter (Nater et al., 2021). Terrestrial predator use of marine subsidies is thought to play a role in the top-down control of tundra food webs (Gauthier et al., 2011; Legagneux et al., 2012) but little empirical support exists for this hypothesis to date. It is also unclear how environmental conditions may influence the indirect effects of marine resources on tundra food webs by affecting terrestrial predator access to these resource subsidies.

In this study, we take an ecosystem-level approach toward evaluating how abiotic and biotic factors influence coastal tundra food web dynamics via direct and indirect pathways in a warming Arctic (Figure 1). We first assessed how environmental conditions affect the abundance of primary terrestrial prey (rodents) and the availability of marine subsidies (seal biomass) to a tundra predator, Arctic foxes, during winter, and the response of the Arctic fox population to these food resources. We then evaluated how the fox population response to rodents and marine subsidies, in turn, influenced the reproductive success of Canada geese (*Branta canadensis interior*) during spring,

while accounting for environmental conditions previously shown to affect goose reproduction (snow persistence, temperature, precipitation). We hypothesized that the availability of Arctic fox foods would be greater under harsh winter conditions, with greater snow persistence positively affecting rodent abundance and longer sea ice duration facilitating fox access to more seals. Nests initiated later have lower rates of success in this system (Johnson-Bice, Warret Rodrigues, et al., 2025), and we hypothesized that snow persistence on the tundra drives goose nesting phenology; we thus predicted geese initiate nesting later during years snow persists longer. We also predicted rodent abundance and marine subsidy availability positively affected fox den occupancy. Our study area lacks annual data on seal abundance, so we used polar bear energetics as a proxy for marine subsidy availability to Arctic foxes, given that both species feed on the same resource and Arctic foxes frequently scavenge seal carrion left from polar bear kills (Roth, 2002; Stirling & Archibald, 1977). Finally, we predicted goose nest success would be lower during years when fox abundance was high, demonstrating an indirect effect of rodents and/or marine subsidies on goose reproduction via the response of a shared predator (foxes). Thus, this study also tests the hypothesis that terrestrial predator use of allochthonous subsidies may contribute to the top-down control of tundra food webs.

METHODS

Study area

We conducted our study in Wapusk National Park in Manitoba, Canada, along the western coast of Hudson Bay (Figure 2; Johnson-Bice et al., 2023; Roth, 2003). Arctic foxes are the main fox species and predators of goose nests in the area (responsible for >80% of nest failures in some years; Reiter & Andersen, 2011; Walter, 1996). Recent research demonstrated Canada geese proactively choose nesting sites in areas less used by foxes, and nesting success is lower in areas most used by foxes (Johnson-Bice, Warret Rodrigues, et al., 2025). We have located 88 natal fox dens within Wapusk (Johnson-Bice et al., 2023), but this study focuses on the closest 22 dens to the goose nesting study area, which we monitor annually by foot in the goose nesting period. Although red fox occurrence in Wapusk has increased in recent years (Moizan et al., 2023), the 22 dens around the goose study area are still primarily occupied by Arctic foxes, and we focus the hypotheses and framing of our study on Arctic foxes.




FIGURE 1 Conceptual model depicting our hypotheses on how winter environmental conditions, food abundance/availability, and predator abundance interactively shape Canada goose nest success through direct and indirect pathways in a coastal tundra ecosystem. We hypothesized environmental conditions affect the availability of winter food resources for Arctic foxes, with greater snow and sea ice duration having positive effects on rodent abundance and seal carrion availability (marine subsidies), respectively. We also hypothesized that greater fox activity/abundance would negatively affect goose nest success, which would demonstrate indirect negative effects of marine resources and rodents on goose reproduction mediated through a shared predator (foxes). We hypothesized environmental conditions play a strong role in goose reproduction, with snow cover affecting the timing of nest incubation, and temperature and precipitation during incubation affecting the availability of resources for breeding geese. Solid black lines represent hypothesized relationships and pathways evaluated in this study. The solid blue lines relating to seal availability represent latent direct relationships that we could not test due to an absence of seal abundance data. Dashed lines represent the hypothesized indirect effects of rodents, seals, and snow conditions on goose reproduction (dashed blue = latent indirect effect). Illustrations by Sean M. Johnson-Bice.

During winter, Arctic fox diets are heavily influenced by rodents (*Dicrostonyx*, *Microtus*) in this region, but they also consume large amounts of marine resources (namely ringed seal biomass), particularly when rodent abundance is low (Dudenhoeffer et al., 2021; Roth, 2002). In this region, marine resources may constitute up to 67% of Arctic fox diets (Roth, 2003). Recent research showed Arctic fox space use on the nearby landfast ice overlaps

significantly with both ringed seal and polar bear space use (Warret Rodrigues et al., 2025), highlighting the intricate relationships between these species in the area. Preliminary results from an ongoing study further indicate Arctic foxes are the most common scavenger of seal carcasses (H.E.L. Gamblin, University of Manitoba, personal communication). Although caribou carrion can be an important winter food for Arctic foxes in some areas,




FIGURE 2 Map of our study area within Wapusk National Park, Manitoba, Canada, along the western coast of Hudson Bay. Right panel shows all 88 known fox dens on the tundra within Wapusk, with the subset of “focal dens” used for the present study in the inset map (orange circles). Lower left panel shows the goose nest study area (dark gray polygon outline) with one year of 2-km goose nest transects depicted (horizontal gray lines), and the 22 focal dens around the goose nest study area.

caribou is a secondary food source in this system, as the local herd migrates tens to hundreds of kilometers inland from the study area during winter (Dudenhoeffer et al., 2021; Lochansky et al., 2024). Arctic fox summer and fall diets are heavily influenced by geese, but goose resources appear to have a minimal effect on Arctic fox reproduction (McDonald et al., 2017).

This study uses individual Canada goose nest data from 2014 to 2024. The most common goose species in our study area is the Canada goose, but a lesser snow goose (*Chen caerulescens caerulescens*) colony historically occurred ~12 km west (Cooke et al., 1995). For simplicity, we limit our analysis to only Canada goose nests, which comprised >95% of goose nests detected.

Data sources

We first assessed the effects of winter food abundance and availability (rodent abundance and marine subsidy availability) on Arctic foxes and the environmental conditions affecting these foods. We estimated rodent abundance (in numbers per hectare) using mark–recapture methods from animals live-trapped in June (full details in McDonald et al., 2017), pooling together both meadow vole (*Microtus pennsylvanicus*) and collared lemming (*Dicrostonyx richardsoni*) captures to estimate total rodent abundance (although, collared lemmings comprise ~87% of captures).

We used polar bear energetics data from the Western Hudson Bay population as a proxy for seal availability (pups and carrion; annual seal abundance data are unavailable). Energetic models of polar bears indicate sea ice availability strongly influences polar bear body condition by controlling access to ice-dependent seals and the length of time bears remain on land, largely fasting, during the ice-free season (Johnson et al., 2020; Pagano et al., 2020; Stirling et al., 1999). Because Arctic foxes and polar bears both feed on seals on the sea ice, we assumed that polar bear energetics, which are influenced by both the abundance and accessibility of prey, adequately represent seal availability to Arctic foxes (i.e., conditions that are good for polar bear hunting success also benefit foxes). This rationale is supported by diet studies showing Arctic foxes frequently consume seal biomass in the region (Dudenhoeffer et al., 2021; Roth, 2002, 2003), the high spatial overlap between Arctic foxes, ringed seals, and polar bears on the nearby landfast ice (Warret Rodrigues et al., 2025), and that regional Arctic fox abundance in the 1980s and 1990s was positively related to polar bear body condition (Roth, 2003). We used energetic estimates of solitary adult female polar bears, as this demographic class has a reduced ability to compete with larger bears for food and annual changes in body condition are thus more likely to reflect environmental conditions (i.e., annual seal availability; Johnson et al., 2020). Energetics data from bears captured during 2014–2023 were used in this study, as only one solitary adult female bear was captured in fall 2024 (avg. 11.6, range: 5–20 in all other years).

We evaluated how environmental conditions affected winter food availability to foxes by assessing the effects of snow cover and sea ice duration on rodent abundance and polar bear energetics, respectively. Our study area lacks detailed snow depth/hardness measurements, so we used satellite imagery to derive estimates of snow cover temporal extent (ordinal date when snow cover disappeared on a per-pixel basis, averaged across the study area) to characterize winter snow conditions for

rodents. Implicit in the relationship between polar bear energetics and sea ice duration is that longer sea ice seasons result in greater polar bear consumption of seals by prolonging the hunting season, allowing bears to store more energy. We used satellite imagery to estimate sea ice break-up date from our study area (ordinal date when sea ice cover in Hudson Bay was <50% for 3 consecutive days). See Appendix S1 for additional details on satellite-derived snow cover data and polar bear energetics data used in this study.

The proportion of dens occupied by foxes during the goose nesting period was used as our index of fox abundance/activity. During 2014–2024 (excluding 2020), we visited 20–22 dens on foot in June to assess fox activity (2014–2016: 20 dens; 2017–2023: 21 dens; 2024: 22 dens). Signs of digging/cleared burrows, fresh prey remains (goose eggshells, goose carcasses, lemming carcasses) and/or fresh scats observed at den sites, coupled with images captured by trail cameras installed on each den, were used to categorize dens as “occupied” or “unoccupied.”

We conducted goose nest surveys using line-transect methods each year between June 8 and 18, 2014–2024 (excluding 2020; see Johnson-Bice, Warret Rodrigues, et al., 2025 for full details of survey methods). Briefly, three to five observers located goose nests using binoculars while walking along fifteen 2-km-long transects. Once a nest was detected, we approached the nest until the goose flushed and used egg flotation methods to estimate the date incubation was initiated, assuming a 28-day incubation period (Reiter & Andersen, 2008). Nests were revisited in late July or early August to assess nest fate; nests were categorized as successful if we observed large pieces of membrane, which correspond to at least one hatched egg (Reiter & Andersen, 2011).

To assess the effects of environmental conditions on nest success, we used daily temperature and total precipitation during the incubation period of each clutch (Juhász et al., 2020). Daily temperature and precipitation data were obtained from Environment Canada’s Churchill weather station.

All surveys and capture and handling procedures were approved by the University of Manitoba Animal Care Committee or Environment and Climate Change Canada’s Western and Northern Animal Care Committee. Research was conducted under permits issued by Parks Canada, the Government of Manitoba, and Environment and Climate Change Canada.

Statistical analyses

We assembled five regression models to evaluate how environmental conditions and trophic interactions

influence the dynamics of this coastal tundra food web (model formulas listed in Appendix S1). Our assembled regression models test the hypothesized relationships between each of the variables identified in our conceptual model (Figure 1). Error distributions of each model were determined based on the values and distribution of the response variable. Although food web dynamics are commonly assessed using structural equation models (SEMs), we were unable to use an SEM here due to the mixture of several different datasets. We note that individual model results would be the same as if we were to fit all models in a piecewise SEM, with the main drawbacks of our approach being that we are unable to perform a directed separation test nor quantify the strength of indirect interactions between consecutive significant paths.

We first assessed (1) the effects of sea ice conditions on the energetics of solitary, adult female polar bears using generalized additive mixed-effects models (GAMMs) using the *mgcv* R package (Wood, 2011). Both storage energy (in megajoules) and energy density (in megajoules per kilogram) are useful metrics for characterizing energetics; storage energy typifies energy available for maintenance, reproduction, and growth, whereas energy density is the amount of energy stored relative to the mass of tissue requiring it for somatic maintenance (Johnson et al., 2020). We fit separate models to both energy density (1a) and storage energy (1b) metrics, with annual sea ice break-up date (ordinal date) as a parametric predictor variable. Storage energy and energy density values were normally distributed, so we used GAMMs with a Gaussian distribution. We hypothesized that the bears' age may also affect bear body condition in a nonlinear manner such that both young and old adults may be in relatively worse body condition due to a reduced ability to compete for and acquire food. We therefore included the estimated age of each bear as a smooth term within each model to account for the possibility that energetics may have nonlinear age-related traits. We included a random intercept term for "Year," but did not include a random intercept for "Bear ID" due to the low number of recaptures in this subset of data (<10%).

We then assessed (2) the effects of snow persistence (ordinal date) on rodent abundance using a generalized linear model with a Tweedie distribution within the *glmmTMB* package (Brooks et al., 2017). The Tweedie distribution can accommodate both zero and positive continuous values (both present in our dataset). Implementation of the Tweedie distribution in *glmmTMB* estimates the power parameter using maximum likelihood as part of the model fitting process.

Next, we evaluated (3) the effects of winter food resources on Arctic foxes using a generalized linear

model with a beta distribution using the *glmmTMB* package (Brooks et al., 2017), which is well suited to model continuous proportional data. The proportion of fox dens "occupied" during the goose nesting period each year was the dependent variable (num. occupied/total num. surveyed), with rodent density and average annual polar bear energy metrics as predictor variables (we again assessed energy density [3a] and storage energy [3b] in separate models since both metrics capture slightly different energetic characteristics).

Other work from our area demonstrated the timing of incubation initiation can predict Canada goose nest success, with nests incubated earlier having higher rates of success (Johnson-Bice, Warret Rodrigues, et al., 2025). Snow cover/duration is often assumed to control the timing of goose incubation, but this relationship has not been evaluated in our system. We therefore assessed (4) the effects of snow persistence on the date geese began incubation using a linear mixed-effects model implemented in the *lme4* R package (Bates et al., 2015). Incubation initiation date (ordinal date) of each clutch was the dependent variable, with snow persistence (ordinal date) as the predictor variable. We included a random intercept term for "Transect ID," which accounts for both the annual and spatial clustering of the nest data. We did not include a random intercept term for "Nest ID" due to the relatively low number of repeated nest sites in our dataset (~18%).

Finally, we evaluated (5) the relationship between fox den occupancy and goose nest success using a binomial generalized linear mixed-effects model with a complementary log-log ("cloglog") link using the *glmmTMB* package (Brooks et al., 2017). Nest success was the dependent variable (1 = successful, 0 = unsuccessful), with the annual proportion of fox dens occupied as a predictor variable. Clutches are found at different stages of incubation and thus vulnerable to predation for different lengths of time (i.e., exposure duration), which can bias clutch survival estimates (Shaffer, 2004). We accounted for the exposure duration of each clutch by including the log of exposure duration as an offset term in the model; exposure duration for successful nests was calculated as the number of days between the survey date and expected hatch date, while exposure duration of failed nests was the midpoint between these two dates (Johnson-Bice, Warret Rodrigues, et al., 2025). We also hypothesized that average daily temperature and total precipitation during incubation may influence goose nest success, and predicted that goose nest success would be positively related to both average temperature and cumulative precipitation, as found in other studies

(e.g., Juhasz et al., 2020). Incubation initiation date was also included to account for its effect on nest success (Johnson-Bice, Warret Rodrigues, et al., 2025). We included a random intercept term for “Transect ID.”

We assessed the fit of all models using residual diagnostic tools and verified the lack of collinearity in each model (VIF values all <3). We verified there were no significant temporal trends of any variables tested in our models during our study period (2014–2024) and that there was no residual temporal autocorrelation in our models. Time series plots of all response variables and plots/results from autocorrelation tests can be found in Appendix S1. All analyses were conducted using R version 4.4.1 (R Core Team, 2024).

RESULTS

Influence of winter environment on Arctic fox foods, and subsequent effects on foxes

Winter environmental conditions affected the abundance and availability of the primary winter food sources for Arctic foxes. Using data from 92 solitary adult female polar bears captured on land during fall 2014–2023 (102 total captures), we found a positive relationship between polar bear energy density (in megajoules per kilogram) and the ordinal date of 50% sea ice breakup (Table 1, Figure 3a). However, we found only weak evidence for a relationship between polar bear storage

TABLE 1 Summary results from the five regression models evaluating the direct and indirect drivers of Canada goose reproductive success in Wapusk National Park, Canada.

Model	Model statistics		CI	Statistic	p value
Variable	Estimate	SE			
1a. Polar bear energy density (deviance explained = 20.9%)					
Sea ice breakup (ordinal date)	$\hat{\beta} = 0.207$	0.095	(0.021, 0.393)	$t = 2.17$	0.032
f_{Age}	edf = 2.172	...	...	$F = 1.85$	0.218
f_{year}	edf = 3.002	...	...	$F = 0.80$	0.087
1b. Polar bear storage energy (deviance explained = 26.6%)					
Sea ice breakup (ordinal date)	$\hat{\beta} = 22.53$	12.94	(−2.83, 47.89)	$t = 1.74$	0.085
f_{Age}	edf = 2.416	...	...	$F = 1.63$	0.214
f_{year}	edf = 4.326	...	...	$F = 1.72$	0.012
2. Rodent abundance (pseudo $R^2 = 0.501$)					
Snow duration (ordinal date)	$\hat{\beta} = 0.069$	0.026	(0.014, 0.121)	$z = 2.67$	0.008
3a. Fox den occupancy rate (pseudo $R^2 = 0.858$)					
Rodent abundance	$\hat{\beta} = 0.580$	0.109	(0.367, 0.794)	$z = 5.33$	<0.0001
Polar bear energy density	$\hat{\beta} = 0.073$	0.019	(0.036, 0.109)	$z = 3.93$	<0.0001
3b. Fox den occupancy rate (pseudo $R^2 = 0.846$)					
Rodent abundance	$\hat{\beta} = 0.525$	0.117	(0.296, 0.755)	$z = 4.49$	<0.0001
Polar bear storage energy	$\hat{\beta} = 0.0006$	0.0002	(0.0003, 0.0009)	$z = 3.68$	0.0002
4. Incubation initiation date (Cond. $R^2 = 0.678$, Marg. $R^2 = 0.459$)					
Snow duration (ordinal date)	$\hat{\beta} = 0.554$	0.041	(0.040, 0.714)	$z = 13.52$	<0.0001
$\sigma_{\text{Transect ID}}$	3.095	0.272	(2.56, 3.65)	...	...
5. Canada goose nest success (Cond. $R^2 = 0.569$, Marg. $R^2 = 0.569$)					
Fox den occupancy rate	$\hat{\beta} = -1.963$	0.847	(−3.623, −0.303)	$z = -2.32$	0.020
Incubation initiation date	$\hat{\beta} = -0.217$	0.018	(−0.253, −0.182)	$z = -12.00$	<0.0001
Avg. temp. during incubation	$\hat{\beta} = 0.184$	0.059	(0.068, 0.301)	$z = 3.10$	0.002
Total precip. during incubation	$\hat{\beta} = 0.014$	0.003	(0.008, 0.019)	$z = 5.16$	<0.0001
$\sigma_{\text{TransectID}}$	<0.001	...	...	...	...

Abbreviation: edf, effective degrees of freedom.




FIGURE 3 Results depicting how winter environmental conditions affect winter food availability for Arctic foxes, which, in turn, appear to respond numerically to greater food resources. Panel (a) shows the estimated effect of sea ice duration on female polar bear energy density (in megajoules per kilogram). Individual points show the annual average values $\pm$ SE. Panel (b) depicts the estimated relationship between snow duration and rodent abundance on the tundra. Panels (c) and (d) show the predicted relationships between female polar bear energy density and rodent density, respectively, on fox den occupancy rates. Points in panels (b–d) depict the raw data used in the models. Ribbons represent $\pm 95\%$ CIs of each estimated relationship. Colors and icons correspond to linkages from Figure 1. Illustrations by Sean M. Johnson-Bice.

energy (in megajoules) and sea ice break-up date (Table 1), suggesting sea ice conditions may affect these energetic metrics in polar bears slightly differently. Age of the bear had no substantial effect on storage energy metrics in either model (Table 1), although younger and older bears tended to have lower energy stores (Appendix S1: Figure S3). Total rodent abundance in June was greater during years when snow persisted longer on the tundra in spring (Table 1, Figure 3b).

Arctic fox den occupancy during the goose nesting period was affected by both rodent abundance and

marine subsidy availability. We found a positive relationship between fox den occupancy in June and both polar bear energy density (Figure 3c) and storage energy (Table 1), which we used as proxies for marine subsidy availability. Corrected Akaike information criterion (AIC_c) values for the models with energy density and energy storage were within 2 (−11.86 vs. −11.18), suggesting the energetic metrics performed equally well in explaining fox den occupancy. Rodent abundance positively affected fox den occupancy (Figure 3d), with similar results between both models tested (Table 1).

Effects of environmental conditions and fox activity on Canada goose reproduction

As expected, both environmental conditions and fox activity appeared to influence Canada goose reproduction ($n = 536$ clutches). We found longer snow persistence on the tundra delayed when geese initiated nest incubation (Table 1, Figure 4a). Fox activity affected Canada goose nest success, as demonstrated by a negative effect of fox den occupancy rate on the probability of individual nest success (Table 1, Figure 4b). As reported elsewhere with this dataset

(Johnson-Bice, Warret Rodrigues, et al., 2025), nests incubated later had lower rates of nest success (Table 1). We also found both total precipitation and average daily temperature during the incubation period positively affected individual goose nest success (Table 1, Figure 4c,d).

DISCUSSION

By compiling data sources related to environmental conditions, the abundance and availability of winter food




FIGURE 4 Results depicting how environmental conditions and fox activity affect Canada goose reproduction. Panel (a) shows the estimated effect of snow duration on incubation initiation date. Individual dots show the annual average values $\pm$ SE, highlighting the relatively low intra-annual variation in the timing of incubation throughout the goose population. Panels (b–d) depict the marginal effects of fox den occupancy, cumulative precipitation during incubation, and average daily temperature during incubation, respectively, on the probability of goose nest success. Dots in panel (b) are the average annual nest success values. Ribbons represent $\pm 95\%$ CIs of each estimated relationship. Predictions in panels (b–d) were conditioned on the median average exposure duration (9 days). Colors and icons correspond to linkages from Figure 1. Illustrations by Sean M. Johnson-Bice.

resources to a top predator, and the response of the predator population to these foods, we provide a comprehensive examination of how goose reproduction is affected by direct and indirect pathways in a coastal tundra food web. We found evidence that environmental conditions influence the abundance and availability of winter foods for Arctic foxes, which in turn negatively affect the reproductive success of Canada geese. Greater Arctic fox den occupancy during years polar bears were in better body condition indicates Arctic foxes showed a numerical response to greater marine resource availability, suggesting marine resources can indirectly alter the reproductive success of a tundra prey species via the subsidization of a mobile predator crossing ecosystem boundaries. We also found environmental conditions play a key role in influencing the susceptibility of goose nests to predation by affecting when geese initiate nesting. Our study provides an ecosystem-level perspective on the dynamics of a tundra food web at the terrestrial–marine interface.

Lemming abundance in the Churchill, Manitoba region has declined significantly over the past century, and our results suggest that climate-driven changes in snow conditions are likely a factor in this decline. Although not all lemming populations are affected by winter snow conditions (Gauthier et al., 2024), our finding that lemming abundance is greater following winters that snow persists longer on the tundra supports studies that demonstrate lemming survival and abundance are related to greater snow depth and higher snow quality (soft basal layer) in some areas (Bilodeau et al., 2013a; Fauteux et al., 2015; Kausrud et al., 2008; Poirier et al., 2023). Longer snow persistence may benefit lemmings by prolonging the length of time snow provides some protection from predators (Bilodeau et al., 2013b), and it may also be an indicator of greater snow depth accumulation. Regardless of the exact mechanism, snow persistence has been declining in our study area since at least 2000 (Appendix S1: Figure S4), suggesting changing snow persistence is likely partially responsible for the long-term decline in lemmings; current rodent abundance peaks at ~2/ha, substantially lower than peak abundance in the 1990s (~12/ha; Roth, 2003) and as far back as the 1930s and 1960s (~40/ha or more; Scott, 1993; Shelford, 1943).

Sea ice availability influences polar bear energetics, a metric we used as an indicator of marine subsidy availability to Arctic foxes, suggesting sea ice conditions also affect marine resource availability to foxes. Lacking annual estimates of seal abundance in our area, we used polar bear energetics as a proxy for marine subsidy availability based on the assumption that environmental conditions that promote polar bear hunting success also

benefit Arctic foxes since they feed on the same ice-dependent prey. The relationships between Arctic foxes, polar bears, and seals on the sea ice are well known, with written reports of foxes scavenging seals killed by polar bears as early as 1928 and of foxes killing seal pups in 1879 (Soper, 1928). Our assumption is also supported by Arctic fox diet studies from our area that show fox consumption of seal biomass (Dudenhoeffer et al., 2021; Roth, 2002, 2003) and high spatial overlap between Arctic foxes and ringed seals on the nearby landfast ice (Warret Rodrigues et al., 2025). Nonetheless, the ability of Arctic foxes and polar bears to access seals clearly depends on sea ice availability. Hudson Bay sea ice duration has declined by ~10 days/decade over the past 40 years (Boonstra et al., 2020), and these degrading sea ice conditions are associated with declines in ringed seal body condition, reproductive output, and population size (Ferguson et al., 2017). Western Hudson Bay polar bears have also exhibited long-term reductions in body condition and energetics (Derocher & Stirling, 1995; Johnson et al., 2020; Stirling et al., 1999), causing demographic changes and a similar population decline (Archer et al., 2025; Lunn et al., 2016). Reduced ringed seal and polar bear abundance suggest that the availability of seal biomass to Arctic foxes has also declined through time. Altogether, this study presents key evidence that changing winter climate has reduced the abundance and availability of the main winter foods for Arctic foxes (rodents and seals), and further supports the hypothesis that the ongoing, long-term decline of this Arctic fox population is largely being caused by climate-driven changes in their winter food supply (Verstege et al., 2023).

Consistent with our hypotheses, environmental conditions also affected Canada goose reproduction, with nest success rates positively related to greater precipitation and warmer temperatures during incubation (Table 1, Figure 4c,d). Greater precipitation is thought to reduce the time females spend off nest to drink water, thereby reducing their susceptibility to predation (Juhasz et al., 2020; Lecomte et al., 2009). Foxes also tend to move slower and be less successful in attacking goose nests in wetter tundra environments (Lecomte et al., 2008), suggesting greater precipitation hinders fox mobility and their ability to depredate nests. Regarding temperature, warmer temperatures are thought to reduce embryo mortality and metabolic costs for incubating females (Dickey et al., 2008). Warmer temperatures may also have a bottom-up effect on geese by increasing food availability, as plant productivity tends to be greater during warmer springs (Kelsey et al., 2021). Snow likely disappears earlier during warm springs as well, allowing geese to initiate nesting earlier (Figure 4a; Lameris et al., 2019; Madsen et al., 2007), which, in turn, leads to greater nest

success (Table 1). Female Canada geese also primarily rely on fat reserves for energy while incubating (Murphy & Boag, 1989; Raveling, 1979), so nesting earlier likely means they deplete less of these energy stores. Environmental conditions during incubation thus appear to influence goose nest success by mediating both top-down and bottom-up processes.

Despite the apparent benefits that warmer springs and earlier snowmelt tend to offer Arctic nesting geese, Canada geese in our system are apparently not adjusting their nesting phenology enough to compensate for earlier springs (Appendix S1: Figure S4). Arctic geese nesting phenology has evolved such that goslings hatch during peak vegetation quality, presumably to maximize growth (Dickey et al., 2008). A mismatch in goose nesting phenology and peak vegetation may therefore ultimately reduce gosling growth and survival (Brook et al., 2015; Nolet et al., 2020). Lameris et al. (2019) found early-breeding barnacle geese (*Branta leucopsis*) had the highest rates of nest success in the high Arctic, whereas geese breeding at intermediate dates had the highest rates of success in the low Arctic, suggesting low-Arctic geese may be more capable of adjusting to a warming climate. However, we have no evidence that Canada geese in our low-Arctic system show greater rates of nest success at intermediate dates (Appendix S1: Figure S6), indicating an ongoing phenological mismatch between goose nesting and the onset of spring. This mismatch may be exacerbated in our study area where long-term grubbing by migrating snow geese has degraded large portions of the tundra and reduced food availability to geese (e.g., Jefferies et al., 2006; Kerbes et al., 1990), and be particularly detrimental to geese during the pre-nesting period by restricting their ability to replenish protein reserves following migration (Baldwin et al., 2022). Phenological mismatch may also help explain why Canada goose abundance has remained stable for decades (Luukkonen & Leafloor, 2021) despite warmer springs (Appendix S1: Figure S5), lower Arctic fox abundance (Verstege et al., 2023), and a recent collapse in the nearby colony of lesser snow geese (Hudson Bay Project, 2024), the main competitor of Canada geese.

Our finding that increasing rodent abundance has an immediate, negative indirect effect on goose reproduction via the numerical response of a shared predator contrasts with other studies from the Arctic. Previous research from our study area found no effect of rodent abundance on Canada goose nest success between 1993 and 2004 (Reiter & Andersen, 2011); however, the authors estimated rodent abundance using population reconstruction methods rather than live-trapping, and Arctic foxes were being lethally removed by managers during the study (Walter, 1996), both of which may have affected their

results. Most studies evaluating the indirect effects of rodents on tundra breeding birds found support for the “alternative prey hypothesis,” whereby prey switching behavior by predators during low lemming years reduces bird nest success (e.g., Bêty et al., 2001, 2002; Juhasz et al., 2020; McKinnon et al., 2013). However, most of these prior studies evaluated rodent–goose–predator interactions in systems with higher average rodent densities and breeding goose colonies numbering tens of thousands of pairs (e.g., Bylot Island). Arctic foxes likely incur high foraging costs when hunting in large goose colonies from the increased vigilance and deterrence that colonies offer (Samelius & Alisauskas, 2001), so foxes may be more reluctant to hunt there when easier prey (rodents) are abundant (supported by lower attack rates on goose nests in colonies during high-rodent years; Bêty et al., 2002). In contrast, much of the foraging costs associated with hunting dispersed-nesting birds likely come from longer search times, due to the lower spatial predictability of these nests compared to nests in colonies (Dulude-de Broin et al., 2023), which may limit egg acquisition rates by foxes. In systems with high rodent density, Arctic foxes may also be able to meet their energetic demands by mostly consuming rodents during high-rodent years (Beardsell et al., 2022), thereby releasing predation pressure on geese. Marolla et al. (2019) reported a positive indirect effect of rodent abundance on lesser white-fronted goose (*Anser erythropus*) reproduction, a dispersed-breeding goose species, but rodent abundance was substantially greater in their study area than ours (peak densities of ~111 rodents/ha [40 rodents per 60 × 60 m grid] vs. peak densities of 2.0/ha in this study). Peak rodent abundance in Wapusk may no longer be sufficient to sustain foxes without supplemental alternative prey, so foxes continue to consume geese and their eggs regardless of rodent abundance; indeed, no relationship between rodent abundance and the proportion of Arctic fox diet comprised of geese was previously documented (McDonald et al., 2017). In other words, rodent abundance may not be high enough in Wapusk to elicit prey-switching behavior by foxes during the goose nesting period, resulting in near-term apparent competition between rodents and geese due to greater encounter rates between foxes and nesting geese when rodent abundance is higher. We suggest that absolute goose and rodent abundance, as well as their spatial dispersion, influences whether the indirect effects of rodents on geese are positive or negative, demonstrating some context dependency in the relationship between tundra geese and rodents.

We provide evidence that marine resources may alter tundra food webs via the population response of a mobile terrestrial predator, a dynamic that is likely widespread throughout the Arctic due to the circumpolar

distributions of Arctic foxes, polar bears, seals, and geese. Arctic fox den occupancy was higher following winters when adult female polar bears were in good body condition (Figure 3c), suggesting foxes respond numerically to greater marine resource availability during winter (Roth, 2003). Arctic foxes in our area tend to disperse in winter during food scarcity (Warret Rodrigues & Roth, 2023), but increased access to marine subsidies evidently leads to greater fox reproduction and/or increases in adult survival or retention, resulting in greater den occupancy rates. Our finding that Arctic fox den occupancy, in turn, was negatively related to goose nest success suggests an indirect link between marine subsidies and tundra food web dynamics. Tundra food webs are strongly regulated by top-down forces during summer, and predator use of allochthonous subsidies was previously hypothesized to strengthen these top-down effects (Gauthier et al., 2011; Legagneux et al., 2012) but had never been supported prior to this study. The different timing in the availability of seals and geese likely drives this interaction and allowed us to parse out these relationships. Seal pups/carrion are available only in winter/early spring, which is when fox mating and reproduction occur, while geese are available only during late spring and summer; any indirect effects of seals on goose reproduction would thus be mediated by the fox population's response to seals in winter. These inferences align well with a study from Fennoscandia that showed reindeer carrion abundance in winter can indirectly alter goose reproduction in spring by subsidizing foxes (Marolla et al., 2019). Our study adds to the literature demonstrating the timing of spatial subsidies can play a role in influencing food web dynamics (Sato et al., 2016).

Although this study suggests that Arctic marine and tundra food webs are linked via the subsidization of a terrestrial predator on the sea ice, the rapid climate-driven degradation of coastal Arctic ecosystems suggests that the strength of these interaction pathways is declining and poised for further disruption. When food is abundant, polar bears preferentially strip the blubber off of seals they kill (Stirling & Archibald, 1977), leaving large amounts of carrion for scavengers like Arctic foxes. However, warming Arctic conditions are associated with an ongoing decline in ringed seal abundance (Ferguson et al., 2017), a main prey for Western Hudson Bay polar bears and Arctic foxes. The combination of fewer seals, coupled with fewer polar bears and longer fasting periods for bears (Archer et al., 2025), likely means fewer seal pups available to predators and less carrion available for scavengers, due to fewer seal carcasses generated and higher rates of carcass utilization by bears (Petherick et al., 2021). Shorter sea ice seasons also imply foxes have access to marine resources for shorter periods of time (Nater et al., 2021). Collectively, these patterns indicate

an ongoing and future weakening of the marine subsidization of Arctic foxes that, along with declining lemming abundance and lower overall goose production likely due in part to a reproductive phenological mismatch (Brook et al., 2015), will continue to have consequences for tundra ecosystems. Our work not only sheds light on the linkage between Arctic tundra and marine food webs but also highlights the importance of continued long-term collaborative research efforts and monitoring of these food webs, as typified by the current study, to better understand how the effects of climate change are reshaping trophic interactions through direct and indirect pathways in a rapidly warming Arctic.

AUTHOR CONTRIBUTIONS

Sean M. Johnson-Bice and James D. Roth led the study conceptualization. Sean M. Johnson-Bice led the data curation, data analysis, data visualization, and writing and editing of the manuscript. Sean M. Johnson-Bice, Evan S. Richardson, and James D. Roth helped to collect the data. All authors contributed to the study design and revision of the manuscript.

ACKNOWLEDGMENTS

We thank Dave Allcorn, Jill Larkin, and the many students and research professionals who helped collect data and curate the long-term datasets used in this study. We also thank two anonymous reviewers for comments that improved this manuscript. Funding was provided by the Natural Sciences and Engineering Research Council of Canada, Natural Resources Canada Polar Continental Shelf Program, Environment and Climate Change Canada, Parks Canada, the University of Manitoba Fieldwork Support Program, the Churchill Northern Studies Centre (CNSC) Northern Research Fund, the Province of Manitoba, Earth Ranger Foundation, Isdell Family Foundation, and Wildlife Media.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

DATA AVAILABILITY STATEMENT

Data and code (Johnson-Bice, Baldwin, et al., 2025) are available in Figshare at <https://doi.org/10.6084/m9.figshare.26069104>.

ORCID

Sean M. Johnson-Bice  <https://orcid.org/0000-0001-7538-131X>

REFERENCES

- Adams, L. G., S. D. Farley, C. A. Stricker, D. J. Demma, G. H. Roffler, D. C. Miller, and R. O. Rye. 2010. "Are Inland

- Wolf-Ungulate Systems Influenced by Marine Subsidies of Pacific Salmon?" *Ecological Applications* 20: 251–262.
- Angerbjörn, A., M. Tannerfeldt, and S. Erlinge. 1999. "Predator-Prey Relationships: Arctic Foxes and Lemmings." *Journal of Animal Ecology* 68: 34–49.
- Archer, L. C., S. N. Atkinson, N. J. Lunn, S. R. Penk, and P. K. Molnár. 2025. "Energetic Constraints Drive the Decline of a Sentinel Polar Bear Population." *Science* 387: 516–521.
- Baldwin, F. B., R. T. Alisauskas, and J. O. Leafloor. 2022. "Dynamics of Pre-Breeding Nutrient Reserves in Subarctic Staging Lesser Snow Geese (*Anser caerulescens caerulescens*) and Ross's Geese (*Anser rossii*): Implications for Reproduction." *Avian Conservation and Ecology* 17: art38.
- Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. "Fitting Linear Mixed-Effects Models Using lme4." *Journal of Statistical Software* 67: 1–48.
- Beardsell, A., D. Gravel, J. Clermont, D. Berteaux, G. Gauthier, and J. Bêty. 2022. "A Mechanistic Model of Functional Response Provides New Insights into Indirect Interactions among Arctic Tundra Prey." *Ecology* 103: e3734.
- Bêty, J., G. Gauthier, J.-F. Giroux, and E. Korpimäki. 2001. "Are Goose Nesting Success and Lemming Cycles Linked? Interplay between Nest Density and Predators." *Oikos* 93: 388–400.
- Bêty, J., G. Gauthier, E. Korpimäki, and J.-F. Giroux. 2002. "Shared Predators and Indirect Trophic Interactions: Lemming Cycles and Arctic-Nesting Geese." *Journal of Animal Ecology* 71: 88–98.
- Bilodeau, F., G. Gauthier, and D. Berteaux. 2013a. "The Effect of Snow Cover on Lemming Population Cycles in the Canadian High Arctic." *Oecologia* 172: 1007–16.
- Bilodeau, F., G. Gauthier, and D. Berteaux. 2013b. "Effect of Snow Cover on the Vulnerability of Lemmings to Mammalian Predators in the Canadian Arctic." *Journal of Mammalogy* 94: 813–19.
- Boonstra, R., K. Bodner, C. Bosson, B. Delehanty, E. S. Richardson, N. J. Lunn, A. E. Derocher, and P. K. Molnár. 2020. "The Stress of Arctic Warming on Polar Bears." *Global Change Biology* 26: 4197–4214.
- Brook, R. W., J. O. Leafloor, K. F. Abraham, and D. C. Douglas. 2015. "Density Dependence and Phenological Mismatch: Consequences for Growth and Survival of Sub-Arctic Nesting Canada Geese." *Avian Conservation and Ecology* 10(1): art1.
- Brooks, M. E., K. Kristensen, K. J. van Benthem, A. Magnusson, C. W. Berg, A. Nielsen, H. J. Skaug, et al. 2017. "glmmTMB Balances Speed and Flexibility among Packages for Zero-Inflated Generalized Linear Mixed Modeling." *The R Journal* 9: 378–400.
- Carbonell Ellgutter, J. A., D. Ehrich, S. T. Killengreen, R. A. Ims, and E. R. Unnsteinsdóttir. 2020. "Dietary Variation in Icelandic Arctic Fox (*Vulpes lagopus*) over a Period of 30 Years Assessed through Stable Isotopes." *Oecologia* 192: 403–414.
- Chevallier, C., G. Gauthier, S. Lai, and D. Berteaux. 2020. "Pulsed Food Resources Affect Reproduction but Not Adult Apparent Survival in Arctic Foxes." *Oecologia* 193: 557–569.
- Cooke, F., D. B. Lank, and R. F. Rockwell. 1995. *The Snow Geese of La Pérouse Bay: Natural Selection in the Wild*. Oxford: Oxford University Press.
- Derocher, A. E., and I. Stirling. 1995. "Temporal Variation in Reproduction and Body Mass of Polar Bears in Western Hudson Bay." *Canadian Journal of Zoology* 73: 1657–65.
- Dickey, M.-H., G. Gauthier, and M.-C. Cadieux. 2008. "Climatic Effects on the Breeding Phenology and Reproductive Success of an Arctic-Nesting Goose Species." *Global Change Biology* 14: 1973–85.
- Duchesne, D., G. Gauthier, and D. Berteaux. 2011. "Habitat Selection, Reproduction and Predation of Wintering Lemmings in the Arctic." *Oecologia* 167: 967–980.
- Dudenhoefter, M., J. D. Roth, L. K. Johnson, and S. D. Petersen. 2021. "Arctic Fox Winter Dietary Response to Damped Lemming Cycles Estimated from Fecal DNA." *Journal of Mammalogy* 102: 1455–65.
- Dulude-de Broin, F., J. Clermont, A. Beardsell, L.-P. Ouellet, P. Legagneux, J. Bêty, and D. Berteaux. 2023. "Predator Home Range Size Mediates Indirect Interactions between Prey Species in an Arctic Vertebrate Community." *Journal of Animal Ecology* 92: 2373–85.
- Ehrich, D., M. Cerezo, A. Y. Rodnikova, N. A. Sokolova, E. Fuglei, V. G. Shtro, and A. A. Sokolov. 2017. "Vole Abundance and Reindeer Carcasses Determine Breeding Activity of Arctic Foxes in Low Arctic Yamal, Russia." *BMC Ecology* 17: 32.
- Eide, N. E., A. Stien, P. Prestrud, N. G. Yoccoz, and E. Fuglei. 2012. "Reproductive Responses to Spatial and Temporal Prey Availability in a Coastal Arctic Fox Population." *Journal of Animal Ecology* 81: 640–48.
- Eriksson, C. E., D. L. Z. Kantek, S. S. Miyazaki, R. G. Morato, M. dos Santos-Filho, J. S. Ruprecht, C. A. Peres, and T. Levi. 2022. "Extensive Aquatic Subsidies Lead to Territorial Breakdown and High Density of an Apex Predator." *Ecology* 103: e03543.
- Fauteux, D., G. Gauthier, and D. Berteaux. 2015. "Seasonal Demography of a Cyclic Lemming Population in the Canadian Arctic." *Journal of Animal Ecology* 84: 1412–22.
- Ferguson, S. H., B. G. Young, D. J. Yurkowski, R. Anderson, C. Willing, and O. Nielsen. 2017. "Demographic, Ecological, and Physiological Responses of Ringed Seals to an Abrupt Decline in Sea Ice Availability." *PeerJ* 5: e2957.
- Gauthier, G., D. Berteaux, J. Bêty, A. Tarroux, J.-F. Therrien, L. McKinnon, P. Legagneux, and M. C. Cadieux. 2011. "The Tundra Food Web of Bylot Island in a Changing Climate and the Role of Exchanges between Ecosystems." *Ecoscience* 18: 223–235.
- Gauthier, G., D. Ehrich, M. Belke-Brea, F. Domine, R. Alisauskas, K. Clark, F. Ecke, et al. 2024. "Taking the Beat of the Arctic: Are Lemming Population Cycles Changing Due to Winter Climate?" *Proceedings of the Royal Society B: Biological Sciences* 291: 20232361.
- Gilg, O., I. Hanski, and B. Sittler. 2003. "Cyclic Dynamics in a Simple Vertebrate Predator-Prey Community." *Science* 302: 866–68.
- Gilg, O., B. Sittler, B. Sabard, A. Hurstel, R. Sané, P. Delattre, and I. Hanski. 2006. "Functional and Numerical Responses of Four Lemming Predators in High Arctic Greenland." *Oikos* 113: 193–216.
- Hansen, B. B., V. Grotan, R. Aanes, B.-E. Sæther, A. Stien, E. Fuglei, R. A. Ims, N. G. Yoccoz, and Å. Ø. Pedersen. 2013. "Climate Events Synchronize the Dynamics of a Resident Vertebrate Community in the High Arctic." *Science* 339: 313–15.
- Holt, R. D. 1977. "Predation, Apparent Competition, and the Structure of Prey Communities." *Theoretical Population Biology* 12: 197–229.

- Hudson Bay Project. 2024. "The Hudson Bay Project: 2024 Annual Progress Report." 48 pp.
- Jefferies, R. L., A. P. Jano, and K. F. Abraham. 2006. "A Biotic Agent Promotes Large-Scale Catastrophic Change in the Coastal Marshes of Hudson Bay." *Journal of Ecology* 94: 234–242.
- Johnson, A. C., J. R. Reimer, N. J. Lunn, I. Stirling, D. McGeachy, and A. E. Derocher. 2020. "Influence of Sea Ice Dynamics on Population Energetics of Western Hudson Bay Polar Bears." *Conservation Physiology* 8: coaa132.
- Johnson-Bice, S., F. B. Baldwin, E. S. Richardson, and J. Roth. 2025. "Data and R Code For: 'Marine Resources Alter Tundra Food Web Dynamics by Subsidizing a Terrestrial Predator on the Sea Ice.'" Figshare. <https://doi.org/10.6084/m9.figshare.26069104.v1>.
- Johnson-Bice, S. M., J. D. Roth, and J. H. Markham. 2023. "A Cosmic View of 'Tundra Gardens': Satellite Imagery Provides a Landscape-Scale Perspective of Arctic Fox Ecosystem Engineering." *Ecosystems* 26: 1670–84.
- Johnson-Bice, S. M., C. Warret Rodrigues, H. E. L. Gamblin, F. B. Baldwin, and J. D. Roth. 2025. "Predator Activity, Proactive Anti-Predator Strategies, and Nesting Phenology Produce a Dynamic Landscape of Risk to Tundra Goose Reproduction." *Journal of Animal Ecology*. <https://doi.org/10.1111/1365-2656.70128>.
- Juhasz, C.-C., B. Shipley, G. Gauthier, D. Berteaux, and N. Lecomte. 2020. "Direct and Indirect Effects of Regional and Local Climatic Factors on Trophic Interactions in the Arctic Tundra." *Journal of Animal Ecology* 89: 704–715.
- Kausrud, K. L., A. Mysterud, H. Steen, J. O. Vik, E. Østbye, B. Cazelles, E. Framstad, et al. 2008. "Linking Climate Change to Lemming Cycles." *Nature* 456: 93–97.
- Kelsey, K. C., S. H. Pedersen, A. J. Leffler, J. O. Sexton, M. Feng, and J. M. Welker. 2021. "Winter Snow and Spring Temperature Have Differential Effects on Vegetation Phenology and Productivity across Arctic Plant Communities." *Global Change Biology* 27: 1572–86.
- Kerbes, R. H., P. M. Kotanen, and R. L. Jefferies. 1990. "Destruction of Wetland Habitats by Lesser Snow Geese: A Keystone Species on the West Coast of Hudson Bay." *Journal of Applied Ecology* 27: 242–258.
- Lai, S., J. Bêty, and D. Berteaux. 2017. "Movement Tactics of a Mobile Predator in a Meta-Ecosystem with Fluctuating Resources: The Arctic Fox in the High Arctic." *Oikos* 126: 937–947.
- Lameris, T. K., M. E. de Jong, M. P. Boom, H. P. van der Jeugd, K. E. Litvin, M. J. J. E. Loonen, B. A. Nolet, and J. Prop. 2019. "Climate Warming May Affect the Optimal Timing of Reproduction for Migratory Geese Differently in the Low and High Arctic." *Oecologia* 191: 1003–14.
- Lecomte, N., V. Careau, G. Gauthier, and J.-F. Giroux. 2008. "Predator Behaviour and Predation Risk in the Heterogeneous Arctic Environment." *Journal of Animal Ecology* 77: 439–447.
- Lecomte, N., G. Gauthier, and J.-F. Giroux. 2009. "A Link between Water Availability and Nesting Success Mediated by Predator–Prey Interactions in the Arctic." *Ecology* 90: 465–475.
- Legagneux, P., G. Gauthier, D. Berteaux, J. Bêty, M. C. Cadieux, F. Bilodeau, E. Bolduc, et al. 2012. "Disentangling Trophic Relationships in a High Arctic Tundra Ecosystem through Food Web Modeling." *Ecology* 93: 1707–16.
- Lochansky, C., M. R. Boudreau, R. Turner, M. Kliewer, M. Webb, D. Clark, and R. K. Brook. 2024. "Temperature Drives Summer Group Size Dynamics of Eastern Migratory Caribou in the Hudson Bay Lowlands of Manitoba." *Canadian Journal of Zoology* 103: 1–8.
- Lundberg, J., and F. Moberg. 2003. "Mobile Link Organisms and Ecosystem Functioning: Implications for Ecosystem Resilience and Management." *Ecosystems* 6: 87–98.
- Lunn, N. J., S. Servanty, E. V. Regehr, S. J. Converse, E. Richardson, and I. Stirling. 2016. "Demography of an Apex Predator at the Edge of Its Range: Impacts of Changing Sea Ice on Polar Bears in Hudson Bay." *Ecological Applications* 26: 1302–20.
- Luukkonen, D., and J. Leafloor. 2021. "A Management Plan for Mississippi Flyway Canada Geese." Reports Prepared by the Mississippi Flyway Council Technical Section Canada Goose Committee, Mississippi Flyway Council, USA.
- Lydersen, C., and I. Gjertz. 1986. "Studies of the Ringed Seal (*Phoca hispida* Schreber 1775) in its Breeding Habitat in Kongsfjorden, Svalbard." *Polar Research* 4: 57–63.
- Madsen, J., M. Tamstorf, M. Klaassen, N. Eide, C. Glahder, F. Rigét, H. Nyegaard, and F. Cottaar. 2007. "Effects of Snow Cover on the Timing and Success of Reproduction in High-Arctic Pink-Footed Geese *Anser brachyrhynchus*." *Polar Biology* 30: 1363–72.
- Marczak, L. B., R. M. Thompson, and J. S. Richardson. 2007. "Meta-Analysis: Trophic Level, Habitat, and Productivity Shape the Food Web Effects of Resource Subsidies." *Ecology* 88: 140–48.
- Marolla, F., T. Aarvak, I. J. Øien, J. P. Mellard, J.-A. Henden, S. Hamel, A. Stien, T. Tveraa, N. G. Yoccoz, and R. A. Ims. 2019. "Assessing the Effect of Predator Control on an Endangered Goose Population Subjected to Predator-Mediated Food Web Dynamics." *Journal of Applied Ecology* 56: 1245–55.
- McDonald, R. S., J. D. Roth, and F. B. Baldwin. 2017. "Goose Persistence in Fall Strongly Influences Arctic Fox Diet, but Not Reproductive Success, in the Southern Arctic." *Polar Research* 36(sup1): 5.
- McInturf, A. G., L. Pollack, L. H. Yang, and O. Spiegel. 2019. "Vectors with Autonomy: What Distinguishes Animal-Mediated Nutrient Transport from Abiotic Vectors?" *Biological Reviews* 94: 1761–73.
- McKinnon, L., D. Berteaux, G. Gauthier, and J. Bêty. 2013. "Predator-Mediated Interactions between Preferred, Alternative and Incidental Prey in the Arctic Tundra." *Oikos* 122: 1042–48.
- Moizan, A., C. Warret Rodrigues, and J. D. Roth. 2023. "Different Selection Criteria May Relax Competition for Denning Sites between Expanding Red Foxes (*Vulpes vulpes*) and Endemic Arctic Foxes (*Vulpes lagopus*) on the Low-Arctic Tundra." *Polar Biology* 46: 199–213.
- Murphy, A. J., and D. A. Boag. 1989. "Body Reserve and Food Use by Incubating Canada Geese." *The Auk* 106: 439–446.
- Nater, C. R., N. E. Eide, Å. Ø. Pedersen, N. G. Yoccoz, and E. Fuglei. 2021. "Contributions from Terrestrial and Marine Resources Stabilize Predator Populations in a Rapidly Changing Climate." *Ecosphere* 12: e03546.
- Nolet, B. A., K. H. T. Schreven, M. P. Boom, and T. K. Lameris. 2020. "Contrasting Effects of the Onset of Spring on

- Reproductive Success of Arctic-Nesting Geese." *The Auk* 137: ukz063.
- Oli, M. K. 2019. "Population Cycles in Voles and Lemmings: State of the Science and Future Directions." *Mammal Review* 49: 226–239.
- Pagano, A. M., T. C. Atwood, G. M. Durner, and T. M. Williams. 2020. "The Seasonal Energetic Landscape of an Apex Marine Carnivore, the Polar Bear." *Ecology* 101: e02959.
- Petherick, A. S., J. D. Reuther, S. J. Shirar, S. L. Anderson, and L. R. G. DeSantis. 2021. "Dietary Ecology of Alaskan Polar Bears (*Ursus maritimus*) through Time and in Response to Arctic Climate Change." *Global Change Biology* 27: 3109–19.
- Poirier, M., G. Gauthier, F. Domine, and D. Fauteux. 2023. "Lemming Winter Habitat: The Quest for Warm and Soft Snow." *Oecologia* 202: 211–225.
- Polis, G. A., W. B. Anderson, and R. D. Holt. 1997. "Toward an Integration of Landscape and Food Web Ecology: The Dynamics of Spatially Subsidized Food Webs." *Annual Review of Ecology and Systematics* 28: 289–316.
- R Core Team. 2024. *R: A Language and Environment for Statistical Computing [Version 4.4.1]*. Vienna: R Foundation for Statistical Computing.
- Rantanen, M., A. Y. Karpechko, A. Lipponen, K. Nordling, O. Hyvärinen, K. Ruosteenoja, T. Vihma, and A. Laaksonen. 2022. "The Arctic Has Warmed Nearly Four Times Faster than the Globe since 1979." *Communications Earth & Environment* 3: 168.
- Raveling, D. G. 1979. "The Annual Cycle of Body Composition of Canada Geese with Special Reference to Control of Reproduction." *The Auk* 96: 234–252.
- Reiter, M. E., and D. E. Andersen. 2008. "Comparison of the Egg Flotation and Egg Candling Techniques for Estimating Incubation Day of Canada Goose Nests." *Journal of Field Ornithology* 79: 429–437.
- Reiter, M. E., and D. E. Andersen. 2011. "Arctic Foxes, Lemmings, and Canada Goose Nest Survival at Cape Churchill, Manitoba." *The Wilson Journal of Ornithology* 123: 266–277.
- Roffler, G. H., C. E. Eriksson, J. M. Allen, and T. Levi. 2023. "Recovery of a Marine Keystone Predator Transforms Terrestrial Predator–Prey Dynamics." *Proceedings of the National Academy of Sciences of the United States of America* 120: e2209037120.
- Rose, M. D., and G. A. Polis. 1998. "The Distribution and Abundance of Coyotes: The Effects of Allochthonous Food Subsidies from the Sea." *Ecology* 79: 998–1007.
- Roth, J. D. 2002. "Temporal Variability in Arctic Fox Diet as Reflected in Stable-Carbon Isotopes; the Importance of Sea Ice." *Oecologia* 133: 70–77.
- Roth, J. D. 2003. "Variability in Marine Resources Affects Arctic Fox Population Dynamics." *Journal of Animal Ecology* 72: 668–676.
- Samelius, G., and R. T. Alisauskas. 2001. "Deterring Arctic Fox Predation: The Role of Parental Nest Attendance by Lesser Snow Geese." *Canadian Journal of Zoology* 79: 861–66.
- Samelius, G., and R. T. Alisauskas. 2017. "Components of Population Growth for Arctic Foxes at a Large Arctic Goose Colony: The Relative Contributions of Adult Survival and Recruitment." *Polar Research* 36(sup1): 6.
- Sato, T., R. W. El-Sabaawi, K. Campbell, T. Ohta, and J. S. Richardson. 2016. "A Test of the Effects of Timing of a Pulsed Resource Subsidy on Stream Ecosystems." *Journal of Animal Ecology* 85: 1136–46.
- Schmidt, N. M., R. A. Ims, T. T. Hoye, O. Gilg, L. H. Hansen, J. Hansen, M. Lund, et al. 2012. "Response of an Arctic Predator Guild to Collapsing Lemming Cycles." *Proceedings of the Royal Society B: Biological Sciences* 279: 4417–22.
- Scott, P. A. 1993. "Relationship between the Onset of Winter and Collared Lemming Abundance at Churchill, Manitoba, Canada: 1932–90." *Arctic* 46: 293–96.
- Shaffer, T. L. 2004. "A Unified Approach to Analyzing Nest Success." *The Auk* 121: 526–540.
- Shelford, V. E. 1943. "The Abundance of the Collared Lemming (*Dicrostonyx groenlandicus* (TR). VAR. *Richardsoni* Mer.) in the Churchill Area, 1929 to 1940." *Ecology* 24: 472–484.
- Smith, T. G. 1976. "Predation of Ringed Seal Pups (*Phoca hispida*) by the Arctic Fox (*Alopex lagopus*)." *Canadian Journal of Zoology* 54: 1610–16.
- Soper, J. D. 1928. *A Faunal Investigation of Southern Baffin Island*. Ottawa: National Museum of Canada, Canada Department of Mines. Bulletin 53.
- Spiller, D. A., J. Piovio-Scott, A. N. Wright, L. H. Yang, G. Takimoto, T. W. Schoener, and T. Iwata. 2010. "Marine Subsidies Have Multiple Effects on Coastal Food Webs." *Ecology* 91: 1424–34.
- Stenseth, N. C. 1999. "Population Cycles in Voles and Lemmings: Density Dependence and Phase Dependence in a Stochastic World." *Oikos* 87: 427–461.
- Stirling, I., and W. R. Archibald. 1977. "Aspects of Predation of Seals by Polar Bears." *Journal of the Fisheries Research Board of Canada* 34: 1126–29.
- Stirling, I., N. J. Lunn, and J. Iacozza. 1999. "Long-Term Trends in the Population Ecology of Polar Bears in Western Hudson Bay in Relation to Climatic Change." *Arctic* 52: 294–306.
- Subalusky, A. L., and D. M. Post. 2019. "Context Dependency of Animal Resource Subsidies." *Biological Reviews* 94: 517–538.
- Tarroux, A., J. Bêty, G. Gauthier, and D. Berteaux. 2012. "The Marine Side of a Terrestrial Carnivore: Intra-Population Variation in Use of Allochthonous Resources by Arctic Foxes." *PLoS One* 7: e42427.
- Therrien, J.-F., G. Gauthier, and J. Bêty. 2011. "An Avian Terrestrial Predator of the Arctic Relies on the Marine Ecosystem during Winter." *Journal of Avian Biology* 42: 363–69.
- Verstege, J. S., S. M. Johnson-Bice, and J. D. Roth. 2023. "Arctic and Red Fox Population Responses to Climate and Cryosphere Changes at the Arctic's Edge." *Oecologia* 202: 589–599.
- Walter, S. E. 1996. *Aspects of Canada Goose Nesting Ecology in Northern Manitoba: Age, Visibility, and Arctic Fox Predation*. Madison, WI: University of Wisconsin-Madison.
- Warret Rodrigues, C., A. E. Derocher, J. D. Roth, D. McGeachy, and N. W. Pilfold. 2025. "Spatial Overlap of Sea Ice-Associated Predators and Prey in Western Hudson Bay." bioRxiv: 2025.2007.2011.664287.
- Warret Rodrigues, C., and J. D. Roth. 2023. "Feast to Famine: Sympatric Predators Respond Differently to Seasonal Prey Scarcity on the Low Arctic Tundra." *Ecology and Evolution* 13: e9951.

Wood, S. N. 2011. “Fast Stable Restricted Maximum Likelihood and Marginal Likelihood Estimation of Semiparametric Generalized Linear Models.” *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 73: 3–36.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Johnson-Bice, Sean M., Frank B. Baldwin, Evan S. Richardson, and James D. Roth. 2025. “Marine Resources Alter Tundra Food Web Dynamics by Subsidizing a Terrestrial Predator on the Sea Ice.” *Ecology* 106(10): e70204. <https://doi.org/10.1002/ecy.70204>