

RESEARCH ARTICLE

Predator activity, proactive anti-predator strategies and nesting phenology produce a dynamic landscape of risk to tundra goose reproduction

Sean M. Johnson-Bice¹  | Chloé Warret Rodrigues¹  | Holly E. L. Gamblin¹  | Frank B. Baldwin² | James D. Roth¹

¹Department of Biological Sciences,
University of Manitoba, Winnipeg,
Manitoba, Canada

²Environment and Climate Change
Canada, Canadian Wildlife Service,
Winnipeg, Manitoba, Canada

Correspondence

Sean M. Johnson-Bice
Email: s.johnsonbice@gmail.com

Funding information

University of Manitoba; National
Geographic Society; Natural Sciences and
Engineering Research Council of Canada;
Churchill Northern Studies Centre;
Natural Resources Canada

Handling Editor: Francesca Cagnacci

Abstract

1. Birds generally rely on proactive anti-predator strategies when selecting nest sites, as they have limited options to adapt to changing levels of risk once incubation begins. Arctic waterfowl often nest colonially as an anti-predator strategy, but dispersed-breeding species may use other proactive strategies, such as nesting in areas perceived to be safer. However, empirical links between spatial patterns of predation risk and nest habitat selection or success are needed to better understand how predator activity shapes Arctic waterfowl reproduction.
2. Using activity data from the main cursorial nest predators, Arctic and red foxes (*Vulpes lagopus*, *Vulpes vulpes*), and aerial predators, we evaluate the influence predator activity has on Canada goose (*Branta canadensis interior*) nest habitat selection and success, and how nesting phenology mediates these effects.
3. We compared habitat selection models fit to (i) goose nest locations and (ii) fox locations obtained from GPS collars and found that geese and foxes displayed nearly opposite patterns of selection for the same landscape traits. Geese selected sites that minimized their probability of encountering foxes while also maximizing their ability to detect foxes. Spatial predictions of fox activity revealed nests located in areas with higher probability of fox use had lower nest success, indicating fox space-use patterns reflect predation risk.
4. Landscape heterogeneity influenced both fox and goose nest habitat selection patterns and, consequently, spatial variation in predation risk and reproductive success. Aerial predators appeared to have a lesser effect on spatial patterns of goose nest habitat selection and success.
5. Spatial patterns of nest success were, however, strongly influenced by nesting phenology. Nests initiated earlier had a greater probability of successfully hatching, but these patterns depended on location; the benefits of nesting early decreased in areas of high fox activity.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2025 The Author(s). *Journal of Animal Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

6. Our study demonstrates the mechanisms by which nesting phenology, predator activity and landscape heterogeneity interact to shape bird reproduction and provides an empirical demonstration of how our understanding of predation risk is enhanced by integrating both spatial and temporal dynamics.

KEYWORDS

anti-predator behaviour, Arctic fox, Canada goose, landscape of fear, movement ecology, nest success, predation risk, predator-prey interactions

1 | INTRODUCTION

Responses to predation risk range from 'reactive responses' to immediate, short-term risk to 'proactive responses' to risk that occur before interacting with predators (Creel, 2018; Creel et al., 2014). Reactive responses typically involve actions that promote the prey's escape from an attacking predator, such as fleeing or exhibiting reciprocal aggressive behaviour (Bêty et al., 2002; Courbin et al., 2016). Proactive responses are decisions that reduce the probability of interacting with predators, or increase the probability prey will survive an expected encounter (e.g. responses in areas of high predator activity). One way that prey proactively mitigate predation risk is by changing their space-use patterns based on perceived patterns of risk (Lima, 2002). These behavioural responses of prey to predation risk underlie the 'landscape of fear' concept, whereby landscape characteristics, predator space use and prey perception of risk interact to generate expected levels of predation risk across space and time (Gaynor et al., 2019; Laundré et al., 2010). While mobile prey may readily adapt their proactive anti-predator responses to changing levels of risk, immobile prey have fewer options once they are anchored to a location.

Predation is a major source of failure for bird nests, and birds often use proactive strategies to manage risk when selecting nest sites. Proactive anti-predator responses require knowledge of, or direct experience with, predation risk and birds rely on numerous information sources to gauge risk (Chalfoun & Schmidt, 2012; Lima, 2009). For example, birds can obtain information on predation risk while moving throughout the breeding grounds (Reed et al., 1999) or gaining information from previous breeding attempts (Hoover, 2003; Schmidt, 2001). They can then incorporate this information into their nest site decision-making, such as by choosing nest sites in areas of low predator activity (Schmidt et al., 2006) or sites that promote other anti-predator responses (e.g. characteristics that enhance predator detection; Amat & Masero, 2004; Keyel et al., 2013). Despite widespread recognition that predation is a strong selective pressure on bird nesting biology, empirical evaluations linking spatial patterns of risk to nest site selection and survival remain sparse (Clermont et al., 2021; Lima, 2009).

In tundra ecosystems, Arctic birds have adapted proactive anti-predator responses to cope with relatively high exposure to predators due to the short vegetation on the tundra. Smaller

Arctic birds can mitigate nest predation by having cryptically coloured nests/eggs or concealing their nests in the short vegetation (e.g. Ims et al., 2019; Liebezeit et al., 2009), but these strategies are harder for larger species like waterfowl. Consequently, Arctic waterfowl commonly nest in large, dense colonies that offer anti-predator benefits like predator swamping (Baldwin et al., 2011; Iles et al., 2013) and mutual vigilance and mobbing responses (Burger, 1981; Samelius & Alisauskas, 2001). However, some species, like the Canada goose (*Branta canadensis interior*), breed more dispersed. Although Canada geese nest in clumps at broader spatial scales, they exhibit strong territoriality and distance themselves from neighbours at smaller scales (~20–50 m; Reiter & Andersen, 2013). This spacing strategy may help Canada geese reduce encounters with predators by dispersing themselves throughout the landscape, while still being close enough to neighbours to obtain information on attacking predators. When available, Canada geese may select nest sites in microhabitat refuges that restrict terrestrial predators' access, such as on small islets or within colonies of other species to presumably exploit the anti-predator benefits these colonies offer (Baldwin et al., 2011; Corbeil-Robitaille et al., 2024). However, when these anti-predator features are limited, geese often select nest sites in open areas that likely promote visual detection of predators (Miller et al., 2007). Canada geese also show reactive anti-predator responses to predators by aggressively defending their nests from detected predators, with increasing aggression closer to the hatch date (Clermont, Réale, Lindsay, & Giroux, 2019).

Predation risk varies across time as well as space, indicating that an integrated spatiotemporal perspective may better characterize risk (Palmer et al., 2022). For Arctic waterfowl, clear temporal patterns in nest success often emerge based on the date waterfowl begin nesting, with usually greater rates of nest survival the earlier nesting is initiated (Baldwin et al., 2011; Ross et al., 2017). Seasonal decline in nest success can be attributed to factors such as the mother's age and experience (older/more experienced birds tend to nest earlier and have greater nest success; Finney & Cooke, 1978; Raveling, 1981; Forslund & Larsson, 1992; Clermont, Réale, & Giroux, 2019) or body condition (birds that nest later tend to be in worse body condition and have lower rates of nest success; Bêty et al., 2003; Descamps et al., 2011). Predators may, in turn, be better able to predate nests incubated by geese

that are younger, less experienced, or in worse body condition. Thus, nesting phenology—which may reflect seasonal variation in factors like the vulnerability of nesting birds—may interact with predator activity to result in variation in spatial patterns of predation risk across the breeding season. In other words, the probability a nest successfully hatches may depend jointly on when the nest is initiated and the perceived (or real) risk of predation at that nest site.

The over-arching objective of this study was to evaluate how predator activity affects nest habitat selection and nest success of a dispersed-nesting goose, and whether these effects are mediated by nesting phenology. To our knowledge, no direct evaluation has assessed spatial patterns of predation risk on Canada goose nesting ecology, much less how the timing of nesting affects these dynamics, obscuring our understanding of the degree to which predation risk acts as a constraining influence and selective pressure on Canada goose reproduction. Our study was conducted in a low-Arctic ecosystem where the main cursorial goose nest predators are Arctic (*Vulpes lagopus*) and red foxes (*Vulpes vulpes*), along with several avian species that are also egg predators, such as American herring gulls (*Larus smithsonianus*) and common ravens (*Corvus corax*). We first evaluated whether predator activity constrains the spatial distribution of Canada goose nests in the low-Arctic tundra by comparing habitat selection preferences from models—incorporating the same predictor variables—fit to (1) goose nest locations and (2) Arctic and red fox locations obtained with GPS collars. We hypothesized that geese select for nest sites in areas with a low probability of fox use to minimize encounters with foxes and therefore predicted that habitat selection patterns would be opposite for goose nest and fox locations. We then evaluated whether spatial patterns of predator activity can predict the reproductive success of geese (i.e. evaluating whether nest success was lower in areas of high predator activity), and how nesting phenology affects nest survival rates across varying levels of expected predation risk. Together, this approach allowed us to evaluate to what degree predator activity acts as a selective pressure on goose reproduction by generating expected spatial variation in predation risk, and how incorporating temporal dynamics into these spatial patterns, characterized by nesting phenology, may improve our understanding of predator-prey dynamics in tundra ecosystems.

2 | MATERIALS AND METHODS

2.1 | Study area

Our study was conducted in Wapusk National Park in northeastern Manitoba, Canada, along the coast of Hudson Bay (Figure 1). The Wapusk tundra contains large swaths of fen, lakes and ponds and elevated beach ridges formed by isostatic rebound that run roughly parallel to the coastline (Ritchie, 1956; Sella et al., 2007). There are 88 known fox dens within Wapusk, nearly all of which are located on beach ridges (Johnson-Bice et al., 2023; Roth, 2003). Within and

around the goose nesting study area (~32 km²), there are 22 known fox dens, with occupancy rates during the incubation season averaging 0.43 (range: 0.24–0.57; Johnson-Bice et al., [In press](#)).

The most common goose species are the lesser snow goose (*Chen caerulescens caerulescens*) and the Canada goose. These two species exhibit somewhat different nesting strategies: lesser snow geese nest in large colonies, while Canada goose nests are more dispersed (generally at least 20–50 m from nearest neighbours; Reiter & Andersen, 2013). The Wapusk lesser snow goose population dramatically increased in abundance from the 1960s through the 2010s, causing well-documented, widespread changes in tundra vegetation throughout the area (e.g. Cargill & Jefferies, 1984; Jano et al., 1998; Jefferies et al., 2006). Snow geese grub for roots and rhizomes during spring, causing vegetative losses and alterations to the soil sediments (Kerbes et al., 1990). These vegetation changes are most pronounced near the La Pérouse Bay snow goose colony ~12 km west of our core study area (which has declined dramatically in abundance over the past several years; Hudson Bay Project, 2024), but parts of our study area have been altered by goose foraging. Approximately 97% of goose clutches detected in our study area were laid by Canada geese, so we excluded snow goose nests here for simplicity.

Arctic and red foxes are the primary cursorial (i.e. ground-based) goose nest predators in the area, along with occasional predation from polar (*Ursus maritimus*) and grizzly (*Ursus arctos*) bears. Though red fox abundance in the region has remained relatively stable, their presence on the tundra has increased in recent years as Arctic fox abundance has declined (Moizan et al., 2023; Verstege et al., 2023). Rodents (namely collared lemmings *Dicrostonyx richardsoni* and meadow voles *Microtus pennsylvanicus*) tend to be the primary prey for foxes in this area (Dudenhoeffer et al., 2021), but geese are an important seasonal secondary prey (McDonald et al., 2017). Herring gulls are the most common aerial nest predator, but other aerial predators like glaucous gulls (*Larus hyperboreus*), parasitic jaegers (*Stercorarius parasiticus*), common ravens, bald eagles (*Haliaeetus leucocephalus*) and sandhill cranes (*Grus canadensis*) are also relatively common.

2.2 | Goose nest surveys and assessing nest fate

We conducted goose nest surveys between June 8 and 18, 2014–2024 (excluding 2020) using line-transect methods. Each year, we located goose nests along fifteen 2-km long transects spaced 1 km apart (seven transects in the west portion of the study area, eight transects in the east, offset by 500 m; Figure 1). The transects were oriented east–west, generally perpendicular to the beach ridges. The northing value of the northernmost transect was randomly generated each year, resulting in slightly different transects to obtain full coverage of the study area across all study years.

During surveys, 3–5 observers walked along each transect and searched for goose nests using binoculars. Once a nest was detected (usually within 300 m of the transect), we slowly approached




FIGURE 1 Study area within Wapusk National Park, Manitoba, Canada, along the western coast of Hudson Bay. Each of the 10 coloured polygons represents the home range of a collared Arctic fox (*Vulpes lagopus*, solid outlines) or red fox (*Vulpes vulpes*, dashed outlines). The black polygon outline denotes the boundary of the goose nest study area, with the 15 transects from 1 year shown. (The large home range in purple is from a non-breeding Arctic fox that roamed throughout the study area.)

the nest until the goose flushed, then recorded the clutch size and nest location using a handheld GPS unit. Nest processing was done as rapidly as possible (usually <2 min) to minimize disturbance. If eggs were present, we recorded float stage, reflecting development age, which we later used to estimate incubation initiation date (hereafter 'incubation date') assuming a 28-day incubation period (Reiter & Andersen, 2008). If eggs were pipped or goslings were present on the nest, we assumed incubation began 28 days prior to the survey date. If a nest contained fresh down but no eggs, we assumed the nest had either been depredated or the eggs had already hatched. We then determined the fate of nests from the presence (hatched) or absence (depredated) of large pieces of membrane (corresponding to at least one hatched egg) in or near the nest (Reiter & Andersen, 2011). Nests were revisited in late July to early August to assess nest fate, although nest fate data were collected on a random subset of nests before 2018. We also recorded the number of aerial predators observed on each transect as an index of aerial predator activity. There was no indication that aerial predator counts varied by survey date (Supporting Information S1).

2.3 | Fox capture, collaring and home range analysis

We captured Arctic and red foxes during 2018–2023 using padded leghold and Tomahawk live traps (for details on captures and collaring see Warret Rodrigues and Roth (2023a) and Warret Rodrigues and Roth (2023b)). Foxes were fitted with GPS collars that recorded locations every 1.5 or 2 h during the goose nesting season (Iridium satellite collars #4170 or 4270, Telonics, Mesa, AZ, USA). Foxes captured during 2022–2023 ($n=5$) were fitted with collars that recorded locations every 15 min (0.25 h) during one 6-h period each day, with the 6-h period changing daily to ensure locations were recorded across the full 24-h diel period. We thinned the locations obtained from these foxes to one location every 1.5 h to match fix rates of the other foxes. For this study, we included only foxes with home ranges near the goose nesting study area (within 6–8 km, Figure 1; $n=10$ foxes, 2 red and 8 Arctic) during the primary nesting period. All capture, handling and nest survey procedures were approved by the University of Manitoba Animal Care Committee (Protocols F17-012, F22-004). Research

permits were obtained from Parks Canada (WAP-2017-25781, WAP-2018-27938, WAP-2021-38241, WAP-2022-41740), Manitoba Wildlife (WB18911, WB20226, WB21856, WB25383, WB25857) and Environment and Climate Change Canada (14-MB-SC002, 21-MB-SC001).

We constructed home ranges for each collared fox to delineate boundaries within which 'available' locations would be generated for the used-available habitat selection analysis. We first retained only GPS locations collected during the goose nesting period (May 15–June 30) and then generated 95% autocorrelated kernel density home ranges for each fox following guidelines from the *ctmm* R package (see Calabrese et al., 2016 for details).

2.4 | Goose nest and fox habitat selection study design

We compared fox and goose nest habitat selection by fitting two separate models incorporating the same landscape variables in a used-available study design. For both models, we randomly generated 100 available locations for every used (observed) location. Fox available locations were generated within each fox's home range. Goose nest available locations were generated within 300 m of the survey transects (~97% of nests found during surveys were located within this threshold). Some nests were used in multiple years, so only the first observation of a nest was included in the habitat selection model. We excluded all nests categorized as 'open water' from the land cover data set (~12%), as these nests were largely located on small islets within lakes/ponds and the spatial resolution of our habitat data precluded evaluating whether/how geese select for islets. Although islets can be refuges from terrestrial predators (Clermont et al., 2021; Corbeil-Robitaille et al., 2024; Duchesne et al., 2021), our data indicated there was likely no substantial difference in nest success (0.57 on islets vs. 0.51 average success).

We used the same four variables for both habitat selection models to allow for direct comparison: land cover type, distance to nearest patch of open water, distance to nearest beach ridge and an index of terrain ruggedness. Since geese should select for areas that reduce the likelihood of nest depredation, we predicted geese select nest sites close to open water, which would reduce the number of directions from which foxes could approach the nests and potentially limit the distance females needed to travel to drink (Lecomte et al., 2009). We also predicted geese would nest in relatively flat areas to aid in visual detection of predators and far from beach ridges, which contain almost all the fox dens in the area and function as wildlife corridors (beach ridges are one of the few dry habitats).

We used a 5-m resolution data set developed for Wapusk National Park as our land cover data (Ponomarenko et al., 2014), reclassified into seven different categories (Table S1): 'Fen', 'Goose-affected', 'Dryas heath', 'Shallow water/sedge fen', 'Unvegetated', 'Shrub/

willow' and 'Open water'. Goose-affected areas were generally 'Fen' cover types that had been altered by goose foraging, typically characterized with sparse vegetation interspersed with mudflats. Since geese nest on land, we excluded 'Open water' areas from the goose nest analyses but retained it for the fox model, as lakes/ponds are often still frozen during parts of the nesting period and foxes can swim there when unfrozen. Land cover type was then extracted for each used/available point. We also calculated the Euclidean distance (in meters) between each point and the nearest open water patch and beach ridge (Supporting Information S1). We used the 30-m resolution FABDEM elevation data set (Hawker et al., 2022) to create a 5 × 5 cell (150 m by 150 m total area) window terrain ruggedness index (TRI) raster (i.e. calculating relative difference in elevation from a central cell with the neighbouring cells within a 5 × 5 grid) using the 'tri' function from the *spatialEco* R package (Evans & Murphy, 2021). Lower TRI values indicate relatively flat areas, whereas higher TRI values indicate relatively uneven ('rugged') areas.

2.5 | Statistical analyses

2.5.1 | Habitat selection models

We compared fox movement and goose nest locations with random (available) locations using binomial generalized linear mixed effects models (GLMMs) with a logit link. Available locations were assigned a weight of 5000 and used locations a weight of 1 (Fieberg et al., 2021). In each model, we log + 1 transformed the continuous variables (distance to water, distance to beach ridge, TRI) due to the data being heavily skewed and having some values of 0. 'Fen' was the reference land cover type (categorical variable) as it is the most common cover type in the study area. For the fox movement model, we included a random intercept for 'fox ID' and random slopes to allow all continuous covariates to vary by 'fox ID' (Muff et al., 2020). For the goose nest model, we included a random intercept term for 'year'. We used spatial (goose nest model) and individual (fox model) blocking cross-validation methods ($n = 5$ folds) to evaluate the fit of each habitat selection model (Boyce et al., 2002; Roberts et al., 2017; see Supporting Information S1 for full details).

We visualized and compared model results by calculating the relative selection strength (RSS) for each continuous variable from each model, where RSS is a quantitative estimate of the relative magnitude of selection for a given location compared to another location(s) (Augar et al., 2017; Fieberg et al., 2021). We calculated RSS scores across a range of values for each covariate relative to the mean value of that covariate from the goose nest data set, with all other variables fixed. This approach allowed us to make direct comparisons of selection strength between the two models under identical conditions. We used the natural log of RSS values for plotting the effects. Log-RSS values > 0 show selection for those conditions relative to the mean, whereas values < 0 show avoidance relative to the mean.

2.5.2 | Nest fate analysis

We tested the relationship between goose nest success and relative probability of fox use at each nest site using a binomial GLMM with a complementary log-log ('cloglog') link (hatched nests=1, failed nests=0). Throughout the study area, we generated estimates of log-RSS values predicted from the fitted fox movement model and then used the estimated log-RSS value from each nest location as a predictor variable in the nest success model. We included incubation date as another predictor variable, plus an interaction term between incubation date and log-RSS value. Including incubation date meant we excluded depredated clutches that were detected during surveys (~11%). However, given the importance of nesting phenology from previous research, we elected to focus our analysis on nests for which we could estimate incubation date. To evaluate how aerial predator activity influenced nest fate, we included the number of aerial predators observed on the transect as another predictor variable (log-transformed). We accounted for the fact that nests were found at different points during incubation using the logistic-exposure method (Shaffer, 2004). Exposure period for successful nests was calculated as the number of days between the date the nest was found and the expected hatch date, whereas the exposure period for failed nests was calculated as the midpoint between those two dates. The log of the exposure period was then included as an offset term in the model (de Zwaan et al., 2022). We included 'year' as a random intercept term but did not include a random intercept term for 'nest ID' (to account for clutches at the same site from multiple years) due to the low number of repeated nest sites in our data (~18%).

We verified each fitted model had no collinearity issues (variance inflation factors were <2 for each variable) and visually inspected model residuals to assess model fits. We evaluated the importance of individual covariates with an $\alpha=0.05$ and whether the 95% confidence intervals overlapped 0. All GLMMs were fit using the *glmmTMB* R package (Brooks et al., 2017), and we estimated and visualized the marginal effect of fox probability of space use on goose nest success using the 'predict_response' function from the *ggeffects* R package (Lüdtke, 2018). All analyses were carried out using R version 4.4.1 (R Core Team, 2024).

3 | RESULTS

During 2014–2024, we detected 801 Canada goose clutches from 632 unique nesting sites. Of the 632 sites, 75 were classified as 'open water' and were subsequently excluded from the habitat selection and nest fate analyses, leaving 557 sites for these analyses. We assessed the nest fate (successful/failed) of 526 clutches from these sites, 428 of which we also estimated incubation date and recorded the number of aerial predators observed during the survey. Most clutches with a known hatch fate but unknown incubation date were clutches that were already depredated when detected during surveys ($n=55$).

We collected 2048 locations from 10 different GPS-collared foxes (range: 78–397 locations per fox) in the goose nesting study area and vicinity. We detected an average of 12.0 avian predators along each transect ($SD=9.5$, range: 1–58).

3.1 | Habitat selection analysis

Overall, Canada geese selected nest sites ($n=557$) in areas that were predicted to be less used by foxes. Geese selected sites that were close to open water ($\beta=-0.241$ [95% confidence intervals: $-0.312, -0.170$]). Although foxes showed no significant response to water ($\beta=0.045$ [$-0.103, 0.193$]), the 95% confidence intervals of the coefficient estimates and log-RSS predictions did not overlap between the two models (Figure 2a). Goose nests tended to be farther from beach ridges ($\beta=0.265$ [$0.194, 0.337$]), whereas foxes selected for areas close to beach ridges ($\beta=-0.134$ [$-0.245, -0.023$]; Figure 2b). Geese selected nest sites located in relatively flat areas (i.e. lower TRI values; $\beta=-0.736$ [$-0.969, -0.504$]), whereas foxes selected for relatively uneven areas ($\beta=0.579$ [$0.282, 0.877$]; Figure 2c).

Foxes and geese also showed different selection preferences for land cover types. Nests were disproportionately located within areas affected by goose foraging (i.e. goose-affected areas; Figure 2d,e). Geese tended to avoid building nests in *Dryas* heath, unvegetated and shrub/willow cover types (Figure 2d,e). Foxes tended to select land cover types in similar frequencies to their availability, with the notable exception of shrub/willow land cover types and open water (Figure 2d). The random intercept/slope terms of 'fox ID' were influential in the fox movement model, whereas the random intercept term of 'year' in the goose nest model had no effect (Table S2).

When comparing spatial predictions of areas likely to be used by foxes and geese, the generally opposite patterns of selection become evident (Figure 3). There was a strong negative correlation between fox movement log-RSS values and goose nest log-RSS values from across the goose nest study area ($r=-0.684$, $p<0.001$, $n=1,262,898$ raster cells [5-m resolution]).

Cross-validation results indicated both habitat selection models were excellent fits to the data. Across testing folds, the fox movement model had an average Spearman's r value of 0.902 (range: 0.802–0.964) and the goose nest model had an average Spearman's r value of 0.940 (range: 0.912–0.963).

3.2 | Nest fate analysis

Annual Canada goose nest success averaged 0.57 ($SD=0.23$, range: 0.18–0.96). Average incubation date across all nests was May 28 (ordinal day 148), with the average annual start of incubation ranging from May 19 (2023) to June 4 (2021).

The probability of nest success was negatively related to predicted fox activity, represented by the estimated log-RSS value from




FIGURE 2 Results from the habitat selection function (HSF) models demonstrating how foxes (blue) and geese (orange) showed nearly opposite patterns of selection for the same landscape features in Wapusk National Park, Manitoba, Canada. Panels (a–c) show the log-relative selection strength values ($\pm 95\%$ confidence intervals [CI]) relative to the mean x-covariate value from the goose nest data (all other covariates were fixed at their mean). Values greater than 0 show selection, whereas values less than 0 show avoidance (model predictions cross 0 at the mean of each x-covariate). Panel (d) shows the mean ($\pm$ std. error) proportion of used and available points from each land cover type across all 10 foxes (light/dark blue) and all 10 years of goose nests. Panel (e) shows the coefficient estimates ($\pm 95\%$ CI) of each land cover type relative to 'Fen', the reference level used in both fox movement and goose nest HSF models.

the fox movement model at each nest location (mapped in Figure 3) ($n=428$ clutches, $\beta=-15.76$ [$-26.64, -4.88$]; Table S3). Incubation date had the greatest relative influence on nest success, with clutches incubated earlier having a greater probability of success ($\beta=-0.192$ [$-0.231, -0.153$]). However, these results depended on nest location, as there was a significant interaction between fox log-RSS value and incubation date ($\beta=0.105$ [$0.030, 0.179$]): the benefits of nesting earlier dissipated in areas of high fox activity (Figure 4). There was a tendency for nest success to be lower in areas with higher counts of aerial predators, but the 95% confidence intervals of the coefficient estimate overlapped zero ($\beta=-0.218$ [$-0.462, 0.027$]).

4 | DISCUSSION

Using movement data from the dominant cursorial nest predators, Arctic and red foxes, we demonstrate that predator activity has a strong effect on the spatial distribution and reproductive success of a dispersed-breeding goose on the Arctic tundra. As expected, habitat selection preferences of Canada geese were nearly opposite those of foxes, suggesting that geese select nest sites in areas that minimize interactions with foxes. Goose nest success was lower, on average, in areas with a greater probability of use by foxes, although these effects were strongly mediated by the date geese started incubation. In other words, the probability that




FIGURE 3 Relative probability of use generated from the fox movement (a, left) and goose nest (b, right) habitat selection function (HSF) models. Predictions from both models were binned into 10 equal-sized quantiles. Areas with a low relative selection strength (RSS) are shown in blue/purple and areas with a high RSS value are shown in orange/yellow. The plots show how foxes generally select to move along beach ridges, whereas geese avoid beach ridges and select for goose-affected areas (darkest blue areas in left panel/white areas in right panel are open water, which were excluded in the goose nest HSF model).




FIGURE 4 Predicted effects of fox activity (fox movement log-relative selection strength values) and nest incubation date on the probability of Canada goose nest success ($n=428$ clutches). The effects of fox activity are shown at three different goose incubation dates (mean incubation date: May 28, ± 1 std. deviation). Ribbons represent $\pm 95\%$ confidence intervals of each estimate. Predictions were conditioned on the mean average exposure duration (9.7 days).

nests successfully hatched depended on both where the nest was located and when incubation began (Figure 5). Our study provides a novel spatiotemporal perspective on how predator activity influences the reproduction of prey in a heterogeneous tundra landscape.

4.1 | Proactive anti-predator nesting decisions in response to predator space use

Goose nest habitat selection patterns likely reflected an anti-predator strategy to minimize predation risk, while also enabling the incubating birds to meet their physiological demands. Nests tended to be located near water, within flat areas and far from beach ridges (Figure 2a–c), landscape traits that likely aid in the detection and/or defence of predators. Foxes often circle goose nests during an attack (Bahr, 1989), so nesting close to water should restrict the number of directions from which foxes can attack (i.e. foxes will not attack nests from within a lake), whereas nesting in flat areas probably enhances the detection of incoming predators (Miller et al., 2007). Similarly, geese may avoid nesting near the elevated beach ridges as these features may obscure their ability to detect predators on the far side, and the ridges are also home to fox dens and used as movement corridors for other wildlife (Zhao et al., 2022). Thus, selection for




FIGURE 5 Predicted spatial patterns of Canada goose nest success within the study area, demonstrating the dynamic landscape of risk to goose reproduction that is produced by nesting phenology and spatial variation in fox activity. The figure depicts the probability of nest success within a portion of the study area for nests that were incubated relatively early (a) or late (b). Variation in the probability of nest success within each panel reflects spatial variation in risk derived from the fox habitat selection model.

certain landscape features by predators may create predictable spatial variation in predation risk that birds can adaptively respond to and use to inform nesting decisions (Thomson et al., 2006). However, some of the observed selection patterns may also reflect goose preferences for other resources, particularly drinking water. Though incubating females typically spend <5% of time away from nests (Aldrich & Raveling, 1983), drinking is a primary reason incubating geese leave their nests (Lecomte et al., 2009). Foxes are more successful at attacking nests when the goose is off nest than when they are incubating (Bêty et al., 2002), so nesting close to water would reduce their vulnerability to predation by limiting the length of time geese spend off nest. We speculate food demands likely play a lesser role in influencing goose nesting decisions, as geese largely rely on body reserves for energy while incubating (Murphy & Boag, 1989) and nesting geese in this study showed strong selection for areas with degraded food resources ('goose-affected areas'). These results indicate birds balance resource acquisition needs and safety when deciding where to build nests, a classic trade-off animals face during habitat selection.

Foxes showed strong selection for the elevated beach ridges, likely reflecting both reproductive and foraging behaviours, as their reproductive dens and primary prey (lemmings) are generally concentrated around these features (Johnson-Bice et al., 2023). We did not parse out fox habitat selection preferences based on behavioural state (e.g. foraging vs. resting), so the results reflect time spent by foxes attending home sites and hunting all summer prey species (rodents, Canada geese, snow geese, etc.). Previous research has

shown that foxes select different habitat features based on their behavioural state (Grenier-Potvin et al., 2021), so it is likely that fox and goose nest habitat selection patterns are more similar when foxes hunt geese (i.e. foxes selecting habitats where nests are located). Indeed, we often expect high spatial overlap between hunting predators and prey when prey are immobile (Sih, 1984). However, rather than focusing on fox habitat selection when hunting, our primary interest in this study was to evaluate goose nesting decisions in response to overall patterns of predator space use. From the prey's perspective, the full temporal budget of predator space use, not only time spent hunting, is likely important to consider when selecting nest sites. Independent of the predator's behavioural state, nesting near areas predators use most would be risky, particularly so if the predators are generalist foragers, like foxes, that can easily switch to hunting if they encounter a nest. The negative correlation between fox and goose habitat selection should not imply foxes devote little time to hunting goose nests, but rather that geese select nest sites in areas foxes use less across all behavioural states. Our study highlights the benefit of considering all aspects of predators' behavioural ecology in interpreting space use by both predators and prey, ultimately enhancing our understanding of this predator-prey relationship.

In support of predictions from the 'landscape of fear' concept (Gaynor et al., 2019; Landré et al., 2010), heterogeneity in the physical landscape appears to play a key role in influencing spatial variation in predation risk, and consequently nest success, by mediating the habitat selection of both nesting birds and their predators. Landscape

features like beach ridges, lakes and wetlands influence fox movement and space-use decisions (Grenier-Potvin et al., 2021), and this heterogeneity also clearly influences goose nest habitat selection (Figure 2). Whether geese select areas less used by foxes because they are relative predator refuges or because they offer better opportunities to fend off predators once an attack is launched, the underlying motivation is to minimize predation risk. Although nest success tended to be lower in areas with higher avian predator activity, avian predators likely have a lesser influence on Canada goose nest habitat selection than foxes. The same landscape features that influence fox movements (topography, beach ridges, wetlands) pose little constraint to avian movements, thereby plausibly resulting in less predictable spatial variation in avian predator activity. If avian predator activity strongly affected nest-site selection we may have expected geese to select sites that offer concealment (Miller et al., 2007), but geese in this study tended to avoid cover types that would offer overhead concealment (i.e. shrub/willow cover types; Figure 2d,e). Prey often adapt anti-predator behaviour more strongly to one predator over others in multi-predator systems (Morosinotto et al., 2010; Relyea, 2003), and our findings suggest that landscape heterogeneity may influence the relative strength of prey responses to multiple predators by differentially affecting predator movement patterns (Lecomte et al., 2008).

Proactive selection of nest sites in areas perceived to be safer implies that birds have obtained information on predation risk and used that information to guide nesting decisions in a way that presumably increases fitness. Habitat selection theory posits that animals disproportionately select habitats they perceive offer greater fitness benefits (Fretwell & Lucas, 1969; Northrup et al., 2022), indicating we should logically expect bird nest success is higher in areas with lower predation risk. The negative relationship observed between goose nest success and fox space use indicates the latter adequately predicts spatial variation in predation risk (Clermont et al., 2021) and thus indicates fitness is likely higher for geese that select nest sites in areas less used by foxes. However, it remains unclear what information geese use to assess risk from foxes. They may obtain information on predator activity during the period between arrival on the breeding grounds and incubation (10+ days in our area; Neufeld, 2021), use information from previous breeding attempts (as evidenced by older geese having higher rates of nest success; Raveling, 1981; Rockwell et al., 1993), or obtain information from conspecific neighbours, which often affect the spatial distribution and success of clustered nests (Reiter & Andersen, 2013). Regardless of the source of information on predation risk used, our study indicates nesting birds can adequately gauge spatial patterns of risk, and we provide a rare demonstration that predator activity patterns can act as a selective pressure on bird reproduction.

4.2 | Spatial patterns of predation risk to nesting birds and the value of integrating time

With predation being a leading cause of nest failure (Lima, 2009), interest in understanding how predation and spatial patterns of

predation risk affect nesting birds has increased recently alongside maturation of the 'landscape of fear' concept. Most studies evaluating how spatial variation in predator activity affects bird reproduction, however, have used relatively coarse indices of predator activity or abundance (e.g. track plates, traps, or other methods that estimate occupancy and relative abundance; Schmidt et al., 2006; Davis et al., 2019). Advancements in biologging technology offer opportunities to evaluate how bird nesting decisions are influenced by predator movements, which can provide different inferences of wildlife-habitat relationships than occupancy-based approaches (Bassing et al., 2023). For instance, GPS-collar data from mammalian mesopredators was used to demonstrate how predator habitat selection influences the success and spatial distribution of nesting bobwhite (*Colinus virginianus*) (Harris & O'Connell, 2023). Our general approach of comparing predator and prey habitat selection, and assessing how predator space use affects prey reproduction, is now common in studies of mammalian predator-prey dynamics. We suggest this approach is generalizable and underused in many avian systems. Greater research effort evaluating both predator movements and bird nesting biology will provide insight into how reliable predator activity (and predator behavioural state) can predict reproductive success, and ultimately, a more mechanistic understanding of how nesting birds assess and respond proactively to perceived spatial variation in predation risk (Clermont et al., 2021).

Despite increasing recognition that spatial patterns of predation risk often change across time periods (Palmer et al., 2022), most studies evaluating spatial or temporal patterns of bird nest success tend to evaluate these factors in isolation. Seasonal decline in bird nest success is well documented across diverse ecosystems due to factors such as seasonal variation in breeding experience or body condition of nesting birds (e.g. Bêty et al., 2003; Descamps et al., 2011; Finney & Cooke, 1978; Raveling, 1981), but intraseasonal changes in predator behaviour or abundance may also alter predation risk patterns. Across the nesting season, predators may increase their foraging activity (Bonnefond et al., 2025; Sperry et al., 2008), show seasonal increases in abundance (e.g. in systems with migratory predators; Lameris et al., 2019), or develop their search image for nests (Husby & Verdal, 2024), all of which may result in temporal variation in predation risk across the landscape. Here, we showed that the probability of nest success for a given nest site varied across the nesting season, where the benefits of nesting early declined in areas most used by foxes (Figure 4). Bird nesting season aligns with the offspring rearing season for foxes, and we speculate that greater provisioning demands as offspring grow may affect fox foraging behaviour (as has been shown for other canids like grey wolves *Canis lupus*; Gable et al., 2023) and ultimately predation pressure on birds throughout the nesting season. To our knowledge, this study is one of the first to empirically show that spatial patterns in predation risk to nesting birds vary temporally (Figure 5), though we doubt this pattern is unique to our system. Studies evaluating how spatial patterns of predation risk vary across the nesting season will help assess how

common these patterns are and provide a better understanding of predation's role in bird reproduction.

Without the ability to move nests in response to changing levels of risk, birds often employ proactive anti-predator responses to guide nesting decisions and maximize their reproductive success. Using movement data from the dominant nest predators in our system, we showed how the landscape and predator movements interact to create predictable spatial patterns of predation risk, which geese in our system seemed able to perceive and respond proactively to by constructing nests in safer areas. These spatial patterns of predation risk, however, were strongly mediated by nesting phenology, indicating this landscape of risk is temporally dynamic. Our study thus provides an empirical demonstration of how our understanding of bird reproduction, and predator-prey ecology in general, is enhanced by integrating both spatial and temporal dynamics.

AUTHOR CONTRIBUTIONS

Sean M. Johnson-Bice led the data curation, data analysis, data visualization and writing and editing of the manuscript. Sean M. Johnson-Bice, Chloé Warret Rodrigues, Holly E.L. Gamblin and James D. Roth helped collect the data. All authors contributed to the conceptualization, study design and revising of the manuscript.

ACKNOWLEDGEMENTS

Funding for this study was provided by the Natural Sciences and Engineering Research Council of Canada, Natural Resources Canada Polar Continental Shelf Program, the National Geographic Society, the University of Manitoba Fieldwork Support Program and the Churchill Northern Studies Centre (CNSC) Northern Research Fund. We thank the many students and personnel that participated in our long-term research and monitoring efforts, and Parks Canada, the CNSC and Manitoba's Wildlife Branch for their logistic and administrative assistance. We also thank D. Gillis, E. Nol, A. Costamagna, and two anonymous reviewers for comments that improved the content of this manuscript.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

Data and R code available from the figshare data repository. <https://doi.org/10.6084/m9.figshare.28843832>. (Johnson-Bice et al., 2025).

ORCID

Sean M. Johnson-Bice  <https://orcid.org/0000-0001-7538-131X>
Chloé Warret Rodrigues  <https://orcid.org/0000-0001-5852-8498>
Holly E. L. Gamblin  <https://orcid.org/0000-0002-7153-534X>

REFERENCES

- Aldrich, T. W., & Raveling, D. G. (1983). Effects of experience and body weight on incubation behavior of Canada geese. *The Auk*, 100, 670–679.

- Amat, J. A., & Masero, J. A. (2004). Predation risk on incubating adults constrains the choice of thermally favourable nest sites in a plover. *Animal Behaviour*, 67, 293–300.
- Avgar, T., Lele, S. R., Keim, J. L., & Boyce, M. S. (2017). Relative selection strength: Quantifying effect size in habitat- and step-selection inference. *Ecology and Evolution*, 7, 5322–5330.
- Bahr, J. (1989). *The hunting ecology of arctic foxes (Alopex lagopus) near Cape Churchill, Manitoba* (M.S. thesis). University of Manitoba.
- Baldwin, F. B., Alisauskas, R. T., & Leafloor, J. O. (2011). Nest survival and density of cackling geese (*Branta hutchinsii*) inside and outside a Ross's goose (*Chen rossii*) Colony. *The Auk*, 128, 404–414.
- Bassing, S. B., DeVivo, M., Ganz, T. R., Kertson, B. N., Prugh, L. R., Roussin, T., Satterfield, L., Windell, R. M., Wirsing, A. J., & Gardner, B. (2023). Are we telling the same story? Comparing inferences made from camera trap and telemetry data for wildlife monitoring. *Ecological Applications*, 33, e2745.
- Bêty, J., Gauthier, G., & Giroux, J.-F. (2003). Body condition, migration, and timing of reproduction in snow geese: A test of the condition-dependent model of optimal clutch size. *The American Naturalist*, 162, 110–121.
- Bêty, J., Gauthier, G., Korpimäki, E., & Giroux, J.-F. (2002). Shared predators and indirect trophic interactions: Lemming cycles and arctic-nesting geese. *Journal of Animal Ecology*, 71, 88–98.
- Bonnefond, L., Pinaud, D., Bollache, L., Schmidt, N. M., Lang, J., Hansen, L. H., Sittler, B., Moreau, J., & Gilg, O. (2025). Intraseasonal variations in the spatial behaviour of an Arctic predator. *Movement Ecology*, 13, 13.
- Boyce, M. S., Vernier, P. R., Nielsen, S. E., & Schmiegelow, F. K. A. (2002). Evaluating resource selection functions. *Ecological Modelling*, 157, 281–300.
- Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H. J., Mächler, M., & Bolker, B. M. (2017). glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *The R Journal*, 9, 378–400.
- Burger, J. (1981). A model for the evolution of mixed-species colonies of Ciconiiformes. *The Quarterly Review of Biology*, 56, 143–167.
- Calabrese, J. M., Fleming, C. H., & Gurarie, E. (2016). ctmm: An R package for analyzing animal relocation data as a continuous-time stochastic process. *Methods in Ecology and Evolution*, 7, 1124–1132.
- Cargill, S. M., & Jefferies, R. L. (1984). The effects of grazing by lesser snow geese on the vegetation of a sub-Arctic salt marsh. *Journal of Applied Ecology*, 21, 669–686.
- Chalfoun, A. D., & Schmidt, K. A. (2012). Adaptive breeding-habitat selection: Is it for the birds? *The Auk*, 129, 589–599.
- Clermont, J., Grenier-Potvin, A., Duchesne, É., Couchoux, C., Dulude-de Broin, F., Beardsell, A., Bêty, J., & Berteaux, D. (2021). The predator activity landscape predicts the anti-predator behavior and distribution of prey in a tundra community. *Ecosphere*, 12, e03858.
- Clermont, J., Réale, D., & Giroux, J.-F. (2019). Similarity in nest defense intensity in Canada goose pairs. *Behavioral Ecology and Sociobiology*, 73, 108.
- Clermont, J., Réale, D., Lindsay, M.-È., & Giroux, J.-F. (2019). Plasticity, state-dependency, and individual consistency in Canada goose nest defense behavior. *Behavioral Ecology and Sociobiology*, 73, 66.
- Corbeil-Robitaille, M. Z., Duchesne, É., Fortier, D., Kinnard, C., & Bêty, J. (2024). Linking geomorphological processes and wildlife microhabitat selection: Nesting birds select refuges generated by permafrost degradation in the Arctic. *Biogeosciences*, 21, 3401–3423.
- Courbin, N., Loveridge, A. J., Macdonald, D. W., Fritz, H., Valeix, M., Makuwe, E. T., & Chamaillé-Jammes, S. (2016). Reactive responses of zebras to lion encounters shape their predator-prey space game at large scale. *Oikos*, 125, 829–838.
- Creel, S. (2018). The control of risk hypothesis: Reactive vs. proactive anti-predator responses and stress-mediated vs. food-mediated costs of response. *Ecology Letters*, 21, 947–956.

- Creel, S., Schuette, P., & Christianson, D. (2014). Effects of predation risk on group size, vigilance, and foraging behavior in an African ungulate community. *Behavioral Ecology*, 25, 773–784.
- Davis, H. T., Long, A. M., Baumgardt, J. A., Campbell, T. A., & Morrison, M. L. (2019). Factors affecting nest success and predator assemblage of breeding birds in semiarid grasslands. *Rangeland Ecology & Management*, 72, 385–395.
- de Zwaan, D. R., Drake, A., Camfield, A. F., MacDonald, E. C., & Martin, K. (2022). The relative influence of cross-seasonal and local weather effects on the breeding success of a migratory songbird. *Journal of Animal Ecology*, 91, 1458–1470.
- Descamps, S., Bêty, J., Love, O. P., & Gilchrist, H. G. (2011). Individual optimization of reproduction in a long-lived migratory bird: A test of the condition-dependent model of laying date and clutch size. *Functional Ecology*, 25, 671–681.
- Duchesne, É., Lamarre, J.-F., Gauthier, G., Berteaux, D., Gravel, D., & Bêty, J. (2021). Variable strength of predator-mediated effects on species occurrence in an arctic terrestrial vertebrate community. *Ecography*, 44, 1236–1248.
- Dudenhoefter, M., Roth, J. D., Johnson, L. K., & Petersen, S. D. (2021). Arctic fox winter dietary response to damped lemming cycles estimated from fecal DNA. *Journal of Mammalogy*, 102, 1455–1465.
- Evans, J. S., & Murphy, M. A. (2021). *spatialEco*. R package version 1.3–6.
- Fieberg, J. R., Signer, J., Smith, B., & Avgar, T. (2021). A 'how-to' guide for interpreting parameters in habitat-selection analyses. *Journal of Animal Ecology*, 90, 1027–1043.
- Finney, G., & Cooke, F. (1978). Reproductive habits in the snow goose: The influence of female age. *The Condor*, 80, 147–158.
- Forslund, P., & Larsson, K. (1992). Age-related reproductive success in the barnacle goose. *Journal of Animal Ecology*, 61, 195–204.
- Fretwell, S. D., & Lucas, H. L. (1969). On territorial behavior and other factors influencing habitat distribution in birds. *Acta Biotheoretica*, 19, 16–36.
- Gable, T. D., Johnson-Bice, S. M., Homkes, A. T., & Bump, J. K. (2023). Differential provisioning roles, prey size, and prey abundance shape the dynamic feeding behavior of gray wolves. *Communications Biology*, 6, 1045.
- Gaynor, K. M., Brown, J. S., Middleton, A. D., Power, M. E., & Brashares, J. S. (2019). Landscapes of fear: Spatial patterns of risk perception and response. *Trends in Ecology & Evolution*, 34, 355–368.
- Grenier-Potvin, A., Clermont, J., Gauthier, G., & Berteaux, D. (2021). Prey and habitat distribution are not enough to explain predator habitat selection: Addressing intraspecific interactions, behavioural state and time. *Movement Ecology*, 9, 12.
- Harris, J. P., & O'Connell, T. J. (2023). Mesopredator predation risk limits northern bobwhite nesting habitat. *Ecosphere*, 14, e4516.
- Hawker, L., Uhe, P., Paulo, L., Sosa, J., Savage, J., Sampson, C., & Neal, J. (2022). A 30 m global map of elevation with forests and buildings removed. *Environmental Research Letters*, 17, 024016.
- Hoover, J. P. (2003). Decision rules for site fidelity in a migratory bird, the prothonotary warbler. *Ecology*, 84, 416–430.
- Hudson Bay Project. (2024). *The Hudson Bay Project: 2024 annual progress report*. pp. 48.
- Husby, M., & Verdøl, T. (2024). Experiments reveal that search image might be responsible for seasonal variation in nest-predation rates. *Diversity*, 16, 356.
- Iles, D. T., Rockwell, R. F., Matulonis, P., Robertson, G. J., Abraham, K. F., Davies, J. C., & Koons, D. N. (2013). Predators, alternative prey and climate influence annual breeding success of a long-lived sea duck. *Journal of Animal Ecology*, 82, 683–693.
- Ims, R. A., Henden, J.-A., Strömeng, M. A., Thingnes, A. V., Garmo, M. J., & Jepsen, J. U. (2019). Arctic greening and bird nest predation risk across tundra ecotones. *Nature Climate Change*, 9, 607–610.
- Jano, A. P., Jefferies, R. L., & Rockwell, R. F. (1998). The detection of vegetational change by multitemporal analysis of LANDSAT data: The effects of goose foraging. *Journal of Ecology*, 86, 93–99.
- Jefferies, R. L., Jano, A. P., & Abraham, K. F. (2006). A biotic agent promotes large-scale catastrophic change in the coastal marshes of Hudson Bay. *Journal of Ecology*, 94, 234–242.
- Johnson-Bice, S. M., Baldwin, F. B., Richardson, E. S., & Roth, J. D. In press. Marine resources alter tundra food web dynamics by subsidizing a terrestrial predator on the sea ice. *Ecology*.
- Johnson-Bice, S. M., Roth, J. D., & Markham, J. H. (2023). A cosmic view of 'tundra gardens': Satellite imagery provides a landscape-scale perspective of Arctic fox ecosystem engineering. *Ecosystems*, 26, 1670–1684.
- Johnson-Bice, S. M., Warret Rodrigues, W., Gamblin, H. E. L., Baldwin, F. B., & Roth, J. D. (2025). Data and R code for: 'Predator activity, proactive anti-predator strategies, and nesting phenology produce a dynamic landscape of risk to tundra goose reproduction'. *figshare*. <https://doi.org/10.6084/m9.figshare.28843832>
- Kerbes, R. H., Kotanen, P. M., & Jefferies, R. L. (1990). Destruction of wetland habitats by lesser snow geese: A keystone species on the west coast of Hudson Bay. *Journal of Applied Ecology*, 27, 242–258.
- Keyel, A. C., Strong, A. M., Perlut, N. G., & Reed, J. M. (2013). Evaluating the roles of visual openness and edge effects on nest-site selection and reproductive success in grassland birds. *The Auk*, 130, 161–170.
- Lameris, T. K., de Jong, M. E., Boom, M. P., van der Jeugd, H. P., Litvin, K. E., Loonen, M. J. J. E., Nolet, B. A., & Prop, J. (2019). Climate warming may affect the optimal timing of reproduction for migratory geese differently in the low and high Arctic. *Oecologia*, 191, 1003–1014.
- Laundré, J. W., Hernández, L., & Ripple, W. J. (2010). The landscape of fear: Ecological implications of being afraid. *The Open Ecology Journal*, 3, 1–7.
- Lecomte, N., Careau, V., Gauthier, G., & Giroux, J.-F. (2008). Predator behaviour and predation risk in the heterogeneous Arctic environment. *Journal of Animal Ecology*, 77, 439–447.
- Lecomte, N., Gauthier, G., & Giroux, J.-F. (2009). A link between water availability and nesting success mediated by predator-prey interactions in the Arctic. *Ecology*, 90, 465–475.
- Liebezeit, J. R., Kendall, S. J., Brown, S., Johnson, C. B., Martin, P., McDonald, T. L., Payer, D. C., Rea, C. L., Streever, B., Wildman, A. M., & Zack, S. (2009). Influence of human development and predators on nest survival of tundra birds, Arctic coastal plain, Alaska. *Ecological Applications*, 19, 1628–1644.
- Lima, S. L. (2002). Putting predators back into behavioral predator-prey interactions. *Trends in Ecology & Evolution*, 17, 70–75.
- Lima, S. L. (2009). Predators and the breeding bird: Behavioral and reproductive flexibility under the risk of predation. *Biological Reviews*, 84, 485–513.
- Lüdtke, D. (2018). ggeffects: Tidy data frames of marginal effects from regression models. *Journal of Open Source Software*, 3, 772.
- McDonald, R. S., Roth, J. D., & Baldwin, F. B. (2017). Goose persistence in fall strongly influences Arctic fox diet, but not reproductive success, in the southern Arctic. *Polar Research*, 36(sup1), 5.
- Miller, D. A., Grand, J. B., Fondell, T. F., & Anthony, R. M. (2007). Optimizing nest survival and female survival: Consequences of nest site selection for Canada geese. *The Condor*, 109, 769–780.
- Moizan, A., Warret Rodrigues, C., & Roth, J. D. (2023). Different selection criteria may relax competition for denning sites between expanding red foxes (*Vulpes vulpes*) and endemic Arctic foxes (*Vulpes lagopus*) on the low-Arctic tundra. *Polar Biology*, 46, 199–213.
- Morosinotto, C., Thomson, R. L., & Korpimäki, E. (2010). Habitat selection as an antipredator behaviour in a multi-predator landscape: All enemies are not equal. *Journal of Animal Ecology*, 79, 327–333.
- Muff, S., Signer, J., & Fieberg, J. R. (2020). Accounting for individual-specific variation in habitat-selection studies: Efficient estimation of mixed-effects models using Bayesian or frequentist computation. *Journal of Animal Ecology*, 89, 80–92.
- Murphy, A. J., & Boag, D. A. (1989). Body reserve and food use by incubating Canada geese. *The Auk*, 106, 439–446.

- Neufeld, L. (2021). *Comparing migration ecology among geographically distinct populations of Canada geese (Branta canadensis) and cackling geese (Branta hutchinsii)* (Master's thesis). University of Manitoba.
- Northrup, J. M., Vander Wal, E., Bonar, M., Fieberg, J., Laforge, M. P., Leclerc, M., Prokopenko, C. M., & Gerber, B. D. (2022). Conceptual and methodological advances in habitat-selection modeling: Guidelines for ecology and evolution. *Ecological Applications*, 32, e02470.
- Palmer, M. S., Gaynor, K. M., Becker, J. A., Abraham, J. O., Mumma, M. A., & Pringle, R. M. (2022). Dynamic landscapes of fear: Understanding spatiotemporal risk. *Trends in Ecology & Evolution*, 37, 911–925.
- Ponomarenko, S., Quirouette, J., Sharma, R., & McLennan, D. (2014). *Ecotype mapping report for Wapusk National Park*. Monitoring and Ecological Information. Natural Resource Conservation.
- R Core Team. (2024). *R: A language and environment for statistical computing [computer software]*. R Foundation for Statistical Computing. <https://www.R-project.org/>
- Raveling, D. G. (1981). Survival, experience, and age in relation to breeding success of Canada geese. *The Journal of Wildlife Management*, 45, 817–829.
- Reed, J. M., Boulonier, T., Danchin, E., & Oring, L. W. (1999). Informed dispersal: Prospecting by birds for breeding sites. In V. Nolan, E. D. Ketterson, & C. F. Thompson (Eds.), *Current ornithology* (pp. 189–259). Springer US.
- Reiter, M. E., & Andersen, D. E. (2008). Comparison of the egg flotation and egg candling techniques for estimating incubation day of Canada goose nests. *Journal of Field Ornithology*, 79, 429–437.
- Reiter, M. E., & Andersen, D. E. (2011). Arctic foxes, lemmings, and Canada goose nest survival at Cape Churchill, Manitoba. *The Wilson Journal of Ornithology*, 123, 266–277.
- Reiter, M. E., & Andersen, D. E. (2013). Evidence of territoriality and species interactions from spatial point-pattern analyses of subarctic-nesting geese. *PLoS One*, 8, e81029.
- Relyea, R. A. (2003). How prey respond to combined predators: A review and an empirical test. *Ecology*, 84, 1827–1839.
- Ritchie, J. C. (1956). The native plants of Churchill, Manitoba, Canada. *Canadian Journal of Botany*, 34, 269–320.
- Roberts, D. R., Bahn, V., Ciuti, S., Boyce, M. S., Elith, J., Guilleri-Aroita, G., Hauenstein, S., Lahoz-Monfort, J. J., Schröder, B., Thuiller, W., Warton, D. I., Wintle, B. A., Hartig, F., & Dormann, C. F. (2017). Cross-validation strategies for data with temporal, spatial, hierarchical, or phylogenetic structure. *Ecography*, 40, 913–929.
- Rockwell, R. F., Cooch, E. G., Thompson, C. B., & Cooke, F. (1993). Age and reproductive success in female lesser snow geese: Experience, senescence and the cost of philopatry. *Journal of Animal Ecology*, 62, 323–333.
- Ross, M. V., Alisauskas, R. T., Douglas, D. C., & Kellett, D. K. (2017). Decadal declines in avian herbivore reproduction: Density-dependent nutrition and phenological mismatch in the Arctic. *Ecology*, 98, 1869–1883.
- Roth, J. D. (2003). Variability in marine resources affects arctic fox population dynamics. *Journal of Animal Ecology*, 72, 668–676.
- Samelius, G., & Alisauskas, R. T. (2001). Deterring arctic fox predation: The role of parental nest attendance by lesser snow geese. *Canadian Journal of Zoology*, 79, 861–866.
- Schmidt, K. A. (2001). Site fidelity in habitats with contrasting levels of nest predation and brood parasitism. *Evolutionary Ecology Research*, 3, 553–565.
- Schmidt, K. A., Ostfeld, R. S., & Smyth, K. N. (2006). Spatial heterogeneity in predator activity, nest survivorship, and nest-site selection in two forest thrushes. *Oecologia*, 148, 22–29.
- Sella, G. F., Stein, S., Dixon, T. H., Craymer, M., James, T. S., Mazzotti, S., & Dokka, R. K. (2007). Observation of glacial isostatic adjustment in “stable” North America with GPS. *Geophysical Research Letters*, 34, L02306.
- Shaffer, T. L. (2004). A unified approach to analyzing Nest success. *The Auk*, 121, 526–540.
- Sih, A. (1984). The behavioral response race between predator and prey. *The American Naturalist*, 123, 143–150.
- Sperry, J. H., Peak, R. G., Cimprich, D. A., & Weatherhead, P. J. (2008). Snake activity affects seasonal variation in nest predation risk for birds. *Journal of Avian Biology*, 39, 379–383.
- Thomson, R. L., Forsman, J. T., Sardà-Palomera, F., & Mönkkönen, M. (2006). Fear factor: Prey habitat selection and its consequences in a predation risk landscape. *Ecography*, 29, 507–514.
- Verstege, J. S., Johnson-Bice, S. M., & Roth, J. D. (2023). Arctic and red fox population responses to climate and cryosphere changes at the Arctic's edge. *Oecologia*, 202, 589–599.
- Warret Rodrigues, C., & Roth, J. D. (2023a). Coexistence of two sympatric predators in a transitional ecosystem under constraining environmental conditions: A perspective from space and habitat use. *Movement Ecology*, 11, 60.
- Warret Rodrigues, C., & Roth, J. D. (2023b). Feast to famine: Sympatric predators respond differently to seasonal prey scarcity on the low Arctic tundra. *Ecology and Evolution*, 13, e9951.
- Zhao, S.-T., Johnson-Bice, S. M., & Roth, J. D. (2022). Foxes engineer hotspots of wildlife activity on the nutrient-limited Arctic tundra. *Global Ecology and Conservation*, 40, e02310.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Table S1. Land cover types used in the fox movement and goose nest habitat selection analyses, reclassified from Ponomarenko et al. (2014).

Figure S1. Relationship between the date that goose nest transects were surveyed (range: June 8–18) and the number of predators observed on the transect, fit with a Loess smoother ($\pm 95\%$ confidence intervals). Each point is a single transect.

Table S2. Results from the fox movement and goose nest habitat selection analyses. Parameters in bold were significant at the $\alpha=0.05$ level (excluding the intercept). σ , random effect standard deviation; LC, land cover type; SE, standard error.

Table S3. Results from the nest fate analysis model. Parameters in bold were significant at the $\alpha=0.05$ level (excluding the intercept). σ , random effect standard deviation; SE, standard error.

How to cite this article: Johnson-Bice, S. M., Warret Rodrigues, C., Gamblin, H. E. L., Baldwin, F. B., & Roth, J. D. (2025). Predator activity, proactive anti-predator strategies and nesting phenology produce a dynamic landscape of risk to tundra goose reproduction. *Journal of Animal Ecology*, 94, 2282–2294. <https://doi.org/10.1111/1365-2656.70128>